



FOCUS ON  
**HEALTH**

## **Physiotherapy, Podiatry and Biokinetics**

**Self-study CPD activity**

**Activity Reference Number**

**D3(25) 2029**

**Topic**

Chronic neck pain

**Article(s)**

Effectiveness of myofascial release for adults with chronic neck pain: a meta-analysis

**Profession(s)**

Physiotherapist

Podiatrist

Biokineticist

Supplementary Physiotherapist

Supplementary Podiatrist

Supplementary Biokineticist

Physiotherapy Assistant

Physiotherapy

This activity is approved for **THREE (3) Clinical** Continuing Education Units (CEU's)

Systematic review

# Effectiveness of myofascial release for adults with chronic neck pain: a meta-analysis



Lea Overmann<sup>a,\*</sup>, Robert Schleip<sup>b</sup>, Dennis Anheyer<sup>a</sup>,  
Johannes Michalak<sup>a</sup>

<sup>a</sup> Department of Psychology, University of Witten-Herdecke, Germany

<sup>b</sup> Department of Sport and Health Sciences, Technical University Munich, Germany

## Abstract

**Background** Every second human will experience a phase of neck pain in their lifetime and a high rate of chronicity exists. Because of the complexity and multiple influencing factors, chronic pain conditions are associated with a long treatment and diagnostic process. This leads to a prolonged healing process and high costs.

**Objective** To evaluate the effect of myofascial release on the variables of pain and range of motion in patients with chronic neck pain.

**Method** Selection criteria were set to create a search algorithm for a systematic search in the databases: PubMed, Google Scholar, EBM Reviews, Medline, CINAHL, PEDro, and Science Direct. The risk of bias and the methodological quality was analyzed with the PEDro scale.

**Result** Ten randomized controlled trials, with 549 participants met the eligibility criteria. The methodological quality was ranked from good to excellent. The myofascial release showed a significant difference in pain ( $p = 0.03$ ), rotation to the right ( $p = 0.05$ ), and lateral flexion to the right ( $p = 0.04$ ), compared to other treatment methods. No significant effect was found for improvements in pressure pain threshold.

**Conclusion** Modest effects are observed in pain reduction, suggesting potential benefits of myofascial release in managing chronic neck pain. Further research with standardized protocols and direct comparisons to established therapies is crucial for a comprehensive understanding of myofascial release efficacy.

## Contribution of the paper

- **Effective pain reduction:** Myofascial release might significantly impact the reduction of neck pain.
- **Uncertain Impact on PPT and ROM:** Outcomes for Pressure-Pain Threshold (PPT) and Range of Motion (ROM) remain inconclusive, requiring further research.
- **Treatment Diversity Complexity:** Diverse treatment comparators pose challenges in isolating MFR effects, emphasizing the need for standardized protocols and caution when interpreting the results.

## What does the meta-analysis add to the current literature.

- **Scientific Gap and Need for Further Research:** The meta-analysis highlights a significant scientific gap in definitive evidence supporting myofascial release treatments for chronic neck pain. The identified shortage of studies underscores the need for further research to establish MFR's efficacy, emphasizing standardized protocols, long-term effects, and direct comparisons with established therapies.
- **Consideration of Treatment Modality Variations:** The analysis contributes to understanding the complexity introduced by variations in treatment modalities. The meta-analysis emphasizes the importance of homogeneous study designs to enhance outcome comparability. This insight is valuable for researchers and clinicians aiming to interpret and apply findings from diverse studies in the field of MFR and chronic neck pain.

© 2023 The Author(s). Published by Elsevier Ltd on behalf of Chartered Society of Physiotherapy. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Keywords:** Myofascial release; Chronic pain; Range of motion; Pressure-pain-threshold; Fascia

\* Corresponding author.

E-mail address: [overmann.lea@gmx.de](mailto:overmann.lea@gmx.de) (L. Overmann).

## Introduction

The prevalence of chronic pain in Western societies is estimated at 8% [1]. Neck pain is a common experience, with a high incidence of chronic cases [2]. Chronic primary pain, as classified by the International Classification of Disease (ICD), persists or recurs for longer than 3 months and affects one or more anatomical regions of the body [3]. Middle-aged females with sedentary lifestyles are particularly affected [4]. The head and neck regions are known for their susceptibility to pain syndromes [5]. A survey conducted in Western societies involving 4839 participants revealed that 46% experienced constant pain lasting over 2 years, and 19% had lost their jobs due to chronic pain conditions [6]. The German Federal Statistical Office highlights the association of chronic pain with prolonged treatment and diagnostic processes, resulting in delayed healing and increased costs [7]. In recent decades, the treatment approach for chronic neck pain has shifted towards interdisciplinary pain management that includes physiotherapy as a first-line treatment [2].

An anatomical structure that is hypothesized, to be strongly related to the intensity of pain is the fascial continuum [9]. Even though it has great versatility, scientific research is still sparse [10]. Trauma, inflammations or infections, and structural imbalance of the body lead to a tightening of the fascial strain. If this condition becomes long-term, the loss of physiological adaptability leads to a lack of flexibility, pain, and limitation in movement [11]. At the structural level, the fascia solidifies, the collagen becomes dense and fibrous, and the elastin loses its resiliency, impacting physiological and biomechanical processes and activating nociceptors [9]. Myofascial release (MFR) influences these fascial alterations and is therefore recommended in the treatment of chronic pain [12]. The purpose is to alleviate pain, restore the optimal length of a muscle, improve function, and relieve tissue tenderness of the myofascial system [13]. MFR involves applying a low-load and long-duration stretch to the barriers in the myofascial tissue, with one therapist providing treatment to a single person at a time [14].

Webb and Rajendran 2016 conducted a meta-analysis, to evaluate the effectiveness of MFR in joint ROM and pain, across various body regions.

The MFR group showed significant differences in all outcome domains [15].

Comparable results were found in a meta-analysis regarding the effectiveness of MFR in chronic low back pain [16]. The thoracolumbar fascia has received the most attention in studies on low back pain, while research on the deep fascia of the neck is limited [17]. Current evidence indicates a strong correlation between the facial system and the generation and sustaining of pain, therefore investigating the effectiveness of MFR specifically for the neck is crucial. The objectives of this meta-analysis are to review the existing literature on chronic neck pain and evaluate the effectiveness of MFR. The research question guiding this analysis is:

"Does MFR have a significant impact on reducing pain and improving cervical spine range of motion in patients with chronic neck pain?".

## Method

The present meta-analysis elaborated on the foundation of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [18].

### *Selection criteria*

The PICO Worksheet was used to create the research question and determine keywords for databases, used to carry out an advanced search. The study's content was centered on the combination of "Myofascial Release" and "Chronic neck pain" with a focus on "Randomized Controlled Trials." Synonyms for these terms included: "Soft tissue massage", "Soft tissue manipulation", "MFR", "MFT", and "Fascia". Further keywords were:

"Indirect myofascial release", "Hands-on myofascial release", "Randomized Controlled Trials as Topic", "Clinical Trial", "Randomized trials", "Pain", "Pain perception", "Pain management", "Nociception". The therapeutic intervention utilized in this study is referred to as passive MFR or "hands-on" MFR. Manual techniques are employed by a therapist without the aid of tools or instruments, such as a foam roller. Throughout passive MFR, the therapist utilizes their hands to apply gentle and sustained pressure to targeted areas of the body, with a specific focus on the fascial system. This tactile approach enables the therapist to discern subtle changes within the fascial layers, allowing for a responsive application to facilitate the release of tension. Randomized controlled trials (RCT) in peer-reviewed journals published in English or German language before May 2022 were included. Studies were analyzed in the case of a control group, that received a placebo treatment or different physiotherapeutic interventions. The outcome measures must be the intensity of the pain (VAS or PPT) and/or ROM of the cervical spine. The statistical analysis had to present the mean (M), standard deviation (SD), p-value, and effect size. The methodological quality was examined with the PEDro scale [19]. The trials had to accomplish a score of six (Risk-of-Bias Rating). Studies that did not report pain and/or ROM were excluded, in addition to a variety of pathologies, for example, craniomandibular dysfunction (CMD), migraine, and tension-type headaches.

### *Literature research*

The literature search was conducted from March 2022 to June 2022, utilizing MeSH terms on the PubMed database to facilitate a comprehensive exploration employing standardized medical vocabulary. The search algorithm applied was as follows: ("Myofascial Release"[MeSH Terms] OR

"Myofascial Release" OR "indirect myofascial release" OR "hands-on myofascial release") AND ("Randomized Controlled Trials as Topic"[Mesh] OR "Randomized Controlled Trial" OR "Clinical Trials as Topic"[Mesh] OR "Clinical Trial" OR "randomized trials" OR "clinical trials") AND ("Pain"[MeSH Terms] OR "Pain" OR "pain perception" OR "pain management" OR nociception) AND Humans [Mesh] AND English[lang]. Subsequently, a secondary search was executed on Google Scholar. However, due to limitations in supporting complex search algorithms akin to those in PubMed, a simplified approach was adopted. Various databases, including EBM Reviews, MEDLINE, Cochrane Library, PEDro, and Science Direct, were incorporated. The screening process initially involved title and abstract assessment, followed by the application of inclusion criteria. Lastly, articles underwent critical evaluation for scientific methodology and risk of bias, as detailed in [Table 1](#). The search was conducted by a single researcher without the use of automated tools.

#### *Data extraction*

Specified variables were extracted from the studies: Year of publication, sample size, type of intervention, follow-up period, primary outcomes, and secondary outcomes (see [Table 2](#)). The following scales and measures were used to determine and evaluate the outcomes:

- 1) **Visual Analog Scale:** The variable pain was evaluated with a VAS scale in which a numerical scale was added. It is a straight line with the endpoints; no pain (1) and pain as bad as it could be (10). The patient must classify the perception of the severity of the pain. The VAS is a valuable instrument to assess subjective pain intensity and changes due to therapy [20].
- 2) **Pressure-Pain-Threshold:** The PPT is measured with the application of a pressure algometer. The sensitivity of the deep muscular tissue of the trapezius and sternocleidomastoid muscle is examined. A steadily increasing pressure is given to the muscle tissue until the patient mentions a painful sensation. Higher PPT values are indicative of better outcomes. It is a reliable tool for the assessment of the effect of a treatment, aimed at reducing pain caused by muscle tenderness [21].
- 3) **Range of motion:** The range of motion (ROM) of the cervical spine refers to the maximum extent of joint movement, which includes active and passive mobility [22]. ROM assessment involves active movements in four directions, including flexion, extension, rotation, and lateral flexion, with measurements typically reported in degrees. Higher degrees of ROM are associated with better clinical outcomes.

#### *Risk-of-bias rating*

The PEDro scale was used for the analysis of the methodological quality. The internal and external validity in

the form of generalizability and applicability were examined in addition to the quality of the statistical information provided. It consists of the following 10 items: Random allocation, concealed allocation, baseline comparability, blind subjects, blind therapists, blind assessors, adequate follow-up, intention-to-treat analysis, between-group comparison, point estimates, and variability. The “eligibility criteria” was included as a requirement but not considered in the total score. The items were rated with “Yes” if they were clearly described. Otherwise, they were interpreted as insufficient and rated with “No”. A maximal score of 10 could be reached; higher scores indicated a lower risk of bias and higher methodological quality. Scores between six and eight were “good” and the scores above were “excellent”. If a study met less than five criteria, it was judged as “poor”. In the validation study by Cashin and McAuley, the inter-rater reliability of the PEDro scale was estimated to be fair to excellent (ICC range 0.53 to 0.91). Despite the potential variability in the reliability coefficients, the PEDro scale is a widely used and accepted tool for evaluating the methodological quality of clinical trials in physiotherapy [19].

#### *Statistical analysis*

On condition that at least two studies were available on a specific outcome, pooled analyses were conducted, using R software version 4.2.2 and the meta-package [23]. M and SD were calculated from median, range, interquartile range, minimum, maximum, and first or third quartile if necessary. At first, the M, sample size, and SD were outlined from the pre-and post-measurements of the MFR and control group. Based on these values, the confidence interval from 95% was calculated, if not described in the outcomes of the study. Furthermore, Cohen’s (d) was analyzed to evaluate the effect size and detect differences in the mean of the outcomes of the interventions [24]. Heterogeneity ( $I^2$ ) was used to identify the diversity of the outcomes of the studies [25]. Additionally, the p-value was used to analyze if a statistical significance between the groups exists [26]. To provide a more adequate accounting for uncertainty when pooling treatment effects from a smaller number of heterogeneous studies, the Hartung-Knapp correction was used in this meta-analysis [27].

## **Results**

Conducting an advanced literature search on the PubMed Database, we initially identified 66 articles. After thoroughly reviewing titles and abstracts, four articles that met the eligibility criteria were selected. The exclusion of 62 articles was based on technique-related factors or articles addressing specific pathologies or general unspecific pain. Google Scholar yielded 1700 results. After removing duplicates and screening titles and abstracts against eligibility

Table 1  
PEDro Scale.

Study	Random allocation	Concealed allocation	Baseline Comparability	Blind subjects	Blind therapist	Blind assessors	Adequate follow-up	Intention to treat analysis	Between-group comparison	Point estimate and variability	Total (0 to 10)
Bakar et al. (2014)	Y	Y	Y	N	N	Y	N	Y	Y	Y	7/10
Brück & Jacobi (2021)	Y	Y	Y	N	N	Y	N	Y	Y	Y	7/10
Celenay & Kay (2016)	Y	Y	Y	Y	N	N	N	Y	Y	Y	7/10
El-Gendy & Lashen et al. (2019)	Y	Y	Y	Y	N	N	N	Y	Y	Y	7/10
Gauns & Gurudut (2018)	Y	Y	Y	N	Y	Y	Y	Y	Y	Y	9/10
Namvar & Olyaei et al. (2016)	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	9/10
Retamal & Sejo et al. (2021)	Y	Y	Y	Y	N	Y	N	Y	Y	Y	8/10
Rodriguez-Fuentes et al. (2016)	Y	Y	Y	Y	N	N	Y	Y	Y	Y	8/10
Rodriguez-Huguet et al. (2018)	Y	Y	Y	N	N	Y	Y	Y	Y	Y	8/10
Rodriguez-Huguet et al. (2020)	Y	Y	Y	N	N	Y	Y	Y	Y	Y	8/10

criteria, six additional RCTs remained for inclusion in the meta-analysis. No compatible articles were identified from other sources. A detailed summary can be seen in the PRISMA-Flowchart (see Fig. 1):

*Study characteristics*

The articles were published between May 2014 and August 2021. All in all, 549 patients with chronic neck pain participated in the studies. The sample sizes varied between 34 and 96 participants. Participants' ages ranged from 18 to 69 with a mean value of 40 years. The gender distribution showed a higher number of females (65.63%). Three studies only assured equal distribution of gender but did not present the exact number [28, 29, 30]. The average duration of the treatment sessions was 23 minutes and varied between 10 to 45 minutes. The intervention and control group received between one to twelve treatment sessions. Some of the following interventions were applied in the control group: stability training, strength training, high-velocity low-amplitude techniques (HVLA), laser therapy, stretching, massage, electrotherapy, and manual therapy. One study included a cross-over of the groups. The first group did not receive any intervention, while MFR was applied in the second group. After two weeks the groups made a cross-over [28]. The studies included pre- and post-measurements in which the primary and secondary outcomes were taken directly before and after the intervention. Three articles provided a follow-up period from six days to one month [30,31,32]. Nine articles provided pain as the primary outcome, and six included the ROM of the cervical spine. Three articles dealt with PPT as the primary outcome [28,33,34], while three articles included PPT as the secondary outcome for the assessment of pain [30,38,37]. The clinical studies were conducted in private practices for physiotherapy or hospitals and healthcare centers and were executed by specialized physicians or physiotherapists.

*Methodological quality*

The total score for the rating of the risk of bias assessment and methodological quality of the ten RCTs ranged between seven to nine points, with a mean of 7.8 (see Table 1). Two studies received nine points and were classified as excellent [28,32]. The other studies were assigned as “good”, as they gained a score between seven and eight points. The item with the lowest rating was the “Blinding of the therapist”. One article met the criterion [32]. The outcomes of the studies were taken directly before and immediately after the intervention. An adequate follow-up period was sufficiently fulfilled in three articles [32,35,36]. The random and concealed allocation were met in all articles. Less than 10% of the participants were lost during the follow-up period. The statistical results of the between-group comparison reported more than one key outcome.

Table 2  
Summary of included studies.

Study	Participants	Intervention	Primary Outcome (Mean/Change from baseline)	Secondary Outcome (Mean/Change from baseline)	Follow-up	Outcome Measures
<b>Bakar et al. (2014)</b>	45 Age = 25-45	Ex = Myofascial Release 1 Treatment Sessions of 20 Min. Con = Massage 1 Treatment Sessions of 20 Min.	PPT right M. Sternocleidomastoid (SCM): Ex: -0.6, Con: -1.01 PPT left M. SCM Ex: -1.12, Con: -1.37	Electromyography biofeedback (EMG-BF) EMG-BF SCM right: Ex: 2.91, Con: -2.04 EMG-BF SCM left: Ex: 2.14, Con: -0.91	Pre- and post-Measurement, No follow-up	Pain = PPT, Function= Muscle relaxation response Follow-up = Pre- and Post-Measurement
<b>Brück &amp; Jacobi (2021)</b>	60 Age = 30-65	Ex = Myofascial Release 2 Treatment Session of 15-20 Min. Con1 = HVLA 2 Treatment Session of 15-20 Min. Con2 = No treatment	VAS: Ex: -2.1, Con1: -2.8, Con2: 0.0 ROM of the cervical spine: Flexion: Ex: 9.5, Con1: 6.7, Con2: 0.1 Extension: Ex: 11.1, Con1: 9.9, Con2: 0.5 Lateral flexion left: Ex: 9.2, Con1: 7.2 Con2: -0.2 Lateral flexion right: Ex: 9.9, Con1: 4.8, Con2: 0.8 Rotation left: Ex: 10.3, Con1: 8.8, Con2: 0.1 Rotation right: Ex: 8.9, Con1: 8.5, Con2: -0.5	Neck Pain and Disability Scale (NPDS) NPDS: Ex: -11.3, Con1: -11.3, Con2: 1.1	Pre- and post-Measurement, No follow-up	Pain = VAS, Function = ROM, Neck Pain and Disability Scale Follow-up = Pre- and Post-Measurement
<b>Celenay, &amp; Kaya (2016)</b>	60 Age = 18-65	Ex = Myofascial Release, Stability Training 12 Treatment Sessions, 3 per week of 40-45Min. Con = Stability Training 12 Treatment Sessions, 3 per week of 40-45Min.	VAS Rest: Ex: -2.52, Con: -2.64 Activity: Ex: -2.43, Con: 3.00 Night: Ex: -3.45, Con: -1.42 PPT M. Trapezius right: Ex: 2.34, Con: 0.38 M. Trapezius left: Ex: 2.75, Con: 0.46	Level of anxiety (SSTAI) State Anxiety: Ex: -6.26, Con: -2.27 Trait Anxiety: Ex: -3.53, Con: -1.44 Quality of life (SF-36) Physical component summary (PCS): Ex: 3.89, Con: 2.99 Mental component summary (MCS): Ex: 5.61, Con: 1.91	Pre- and post-Measurement, No follow-up	Pain = VAS, PPT, Level of Anxiety = SSTAI, Quality of Life = SF-36 Follow-up = Pre- and Post-Measurement
<b>El-Genidy &amp; Lasheen et al. (2019)</b>	60 Age = 18-40	Ex = Myofascial Release, Stretching, Strength training 12 Treatment Sessions, 3 per week Con1 = Electrotherapy (Low level laser, ultrasound), Stretching, Strength	VAS: Ex: -3.25, Con1: -3.15, Con2: -1.55 ROM of the cervical spine: Flexion: Ex: 12.75, Con1: 12.95, Con2: 2.65 Extension: Ex: 13.1, Con1: 11.85, Con2: 2.55 Lateral flexion right: Ex:	Neck Disability index (NDI) Ex: -9.23, Con1: -9.3, Con2: -2.9	Pre- and post-Measurement, No follow-up	Pain = VAS, Function = ROM, Neck Disability Index (NDI) Follow-up = Pre- and Post-Measurement

Table 2 (Continued)

Study	Participants	Intervention	Primary Outcome (Mean/Change from baseline)	Secondary Outcome (Mean/Change from baseline)	Follow-up	Outcome Measures
<b>Gauns &amp; Gurudut (2018)</b>	40 Age = 20-50 34	training 12 Treatment Sessions, 3 per week	10.6, Con1: 11.65, Con2: 4.8 Lateral flexion left: Ex: 10.05, Con1: 9.00, Con2: 4.65			
		Con2 = Stretching, Strength training 12 Treatment Sessions, 3 per week	Rotation right Ex: 14.0, Con1: 16.0, Con2: 5.25 Rotation left: Ex: 10.5, Con1: 11.5, Con2: 3.0			
<b>Namvar &amp; Olyaei et al. (2016)</b>	Age = 18-55	Ex = Myofascial Release 15 Min, Conventional Treatment: Hot moist pack 20 Min. TENS 15Min. Stretching, strengthening 15Min.	ROM of the cervical spine: Lateral flexion: Ex: 15.11, Con: 4.32 Rotation: Ex: 17.67, Con: 4.37 VAS: Ex: - 3.41, Con: 0.11 PPT M. Trapezius right: Ex: - 5.56, Con: 0.23	Disability of arm shoulder and hand questionnaire (DASH- Score): Ex: 25.06, Con: 4.37 Extension Power: Ex: - 2.57, Con: - 0.23 NDI: Ex: - 24, Con: - 2.6	Pre- and post- Measurement, 6 days Pre- and post- Measurement, No follow-up	Function = ROM, DASH- Score Follow-up = Pre- and Post- Measurement + 6 days Pain = VAS, PPT, Function = Extension Power, NDI Follow-up = Pre- and Post- Measurement
		Con = Conventional treatment 1 Treatment Session of 50 Min. Ex = Myofascial Release 4 Treatment Sessions of 20 Min. Con = No Treatment				
<b>Retamal &amp; Sejjo et al. (2021)</b>	96 Age = 18-40	Ex = Myofascial Release 2 Treatment Sessions of 10 Min. Con1 = Instrumental Myofascial Release 2 Treatment Sessions of 10 Min. Con2 = Instrumental Myofascial Release, Upper Cervical Manipulation 2 Treatment Sessions of 10 Min.	ROM of the cervical spine Flexion: Ex: 2.52, Con1: 1.00, Con2: 2.12 Extension: Ex: 1.87, Con1: 1.00, Con2: 2.21 Rotation right: Ex: 2.9, Con1: 2.97, Con2: 2.11 Rotation left: Ex: 1.84, Con1: 2.83, Con2: 2.68 Lateral flexion right: Ex: 1.13, Con1: 0.9, Con2: 1.17 Lateral flexion left: Ex: 1.26, Con1: - 0.47, Con2: 1.95 VAS (active) Flexion: Ex: - 0.27, Con1: 0.15, Con2: - 0.68 Extension: Ex: - 0.75, Con1: 0.19, Con2: - 1.09	PPT Suboccipital right: Ex: 0.18, Con1: 0.20, Con2: 0.17 Suboccipital left: Ex: 0.22, Con1: 0.23, Con2: 0.16 M. Trapezius right: Ex: 0.08, Con1: 0.15, G3: 0.00 M. Trapezius left: Ex: 0.08, Con1: 0.09, Con2: 0.00 NDI: Ex: - 2.52, Con1: - 2.33, Con2: - 3.47	Pre- and post- Measurement, 2 weeks	Pain= VAS, PPT, Function = ROM, NDI Follow-up = Pre- and Post- Measurement + 2 weeks

Table 2 (Continued)

Study	Participants	Intervention	Primary Outcome (Mean/Change from baseline)	Secondary Outcome (Mean/Change from baseline)	Follow-up	Outcome Measures
<b>Rodríguez-Fuentes et al. (2016)</b>	59 Age = 18-69	Ex = Myofascial Release 10 Treatment Sessions of 15 Min Con = Manual Therapy 10 Treatment Sessions of 15 Min	Rotation right: Ex: -0.58, Con1: -0.03, Con2: -0.63 Rotation left: Ex: -0.52, Con1: -0.22, Con2: -0.83 Lateral flexion right: Ex: -0.78, Con1: -0.54, Con2: -1.14 Lateral flexion left: Ex: -0.06, Con1: -0.45, Con2: -1.19 VAS: Ex: -1.5, Con: -3.62 ROM of cervical spine Flexion: Ex:3.3, Con: 8.14 Extension: Ex: 4.24, Con: 7.41 Lateral Flexion right: Ex: 2.97, Con: 6.24 Lateral Flexion left: Ex: 4.07, Con: 7.17 Rotation right: Ex: 4.14, Con: 9.17 Rotation left: Ex: 6.17, Con: 11.48 VAS: Ex: -4.75, Con: -3.76	SF-36 PSC: Ex: 1.99, Con: 6.41 MSC: Ex: 1.43, Con: 11.52 Costovertebral (C-V) Angle: Ex: 2.83, Con: 5.0 NDI: Ex: -4.65, Con: -11.53	Pre- and post-Measurement, 5 weeks	Pain = VAS, Function = ROM, NDI, Craniovertebral Angle, Quality of Life = SF-36, Follow-up = Pre- and Post-Measurement + 5 weeks
<b>Rodríguez-Huguet et al. (2018)</b>	41 Age = 20-60	Ex = Myofascial Release 5 Treatment Sessions of 45Min. Con = Multimodal Physiotherapy: Ultrasound 10 Min. TENS 20 Min. Massage 20Min. 5 Treatment Sessions	PPT Suboccipital right: Ex: 0.76, Con: 0.36 Suboccipital left: Ex: 0.63, Con: 0.35 M. Trapezius right: Ex: 0.73, Con: 0.35 M. Trapezius left: Ex: 0.68, Con: 0.43	Pre- and post-Measurement, 4 weeks	Pain = VAS, PPT Follow-up = Pre- and Post-Measurement + 4 weeks	

Table 2 (Continued)

Study	Participants	Intervention	Primary Outcome (Mean/Change from baseline)	Secondary Outcome (Mean/Change from baseline)	Follow-up	Outcome Measures
<b>Rodriguez-Huguet et al. (2020)</b>	54 Age = 20-60	Ex = Myofascial Release 5 Treatment Sessions of 45Min. Con = Multimodal Physiotherapy: Ultrasound 10 Min. TENS 20 Min. Massage 20Min. 10 Treatment Sessions	VAS: Ex: -4.82, Con: -3.78 ROM of cervical spine Flexion: Ex: 11.26, Con: 12.11 Extension: Ex: 11.70, Con: 12.85 Lateral flexion left: Ex: 6.48, Con: 7.00 Lateral flexion right: Ex: 10.85, Con: 10.33 Rotation right: Ex: 8.63, Con: 5.26 Rotation left: Ex: 9.15, Con: 5.33	PPT Suboccipital left: Ex: 0.62, G2: 0.41 Suboccipital right: Ex: 0.72, Con: 0.41 Thoracic left: Ex: 0.63, Con: 0.49 Thoracic right: Ex: 0.72, Con: 0.41	Pre- and post-Measurement, 4 weeks	Pain = VAS, PPT Function = ROM Follow-up = Pre- and Post-Measurement +4 weeks

Ex=Experimental Group, Con=Control Group, Con1=Control Group 1; Con2=Control Group 2

*Effects on the outcome domains*

*Pain*

In our examination of pain outcomes extracted from seven RCTs involving 368 participants, a significant difference emerged ( $d = -1.15$ , 95% CI = [-2.11 to 0.2],  $p = 0.03$ ) with a calculated mean difference (MD) of -1.15 on the pain rating scale (see Fig. 2). A sensitivity analysis was conducted, excluding the Namvar et al. study, which uniquely compared myofascial release with no treatment. Despite the exclusion of this study, the results retained statistical significance ( $p = 0.04$ ) and exhibited a substantial effect size ( $d = 0.86$ , 95% CI = [-1.64 to -0.09]), underscoring the robustness of our findings.

*Pressure pain threshold*

The PPT was measured at the trapezius and the suboccipital areas on the left and the right side. The results of five RCTs with 285 participants were analyzed. A medium overall effect was shown at the left trapezius, 230 participants were included ( $d = 0.59$ , 95%CI = [-0.24 to 1.41],  $p = 0.12$ ). The PPT of the right trapezius was evaluated with the data of 254 participants and represented in four RCTs. There was a medium overall effect size with a non-significant difference ( $d = 0.50$ , 95%CI = [ 0.48 to 1.47],  $p = 0.20$ ). Three studies manifest a significant difference on both sides. Four RCTs were included for the evaluation of the PPT of the suboccipital area (N=205). There was no overall effect and significant difference regarding the right ( $d = 0.17$ , 95%CI = [-0.84 to 1.19],  $p = 0.62$ ) and left ( $d = 0.11$ , 95%CI = [-0.66 to 0.88],  $p = 0.68$ ) suboccipital area (see Fig. 3).

*Range of motion*

Five RCTs provided results including rotation and lateral flexion to the right. The outcomes of 318 participants were included in the evaluation. Lateral flexion to the right showed a significant difference between the groups and a medium effect size ( $d = 0.68$ , 95%CI = [0.04 to 1.32],  $p = 0.04$ ) as well as the rotation to the right ( $d = 0.52$ , 95%CI = [-0.01 to 1.05],  $p = 0.05$ ). Four RCTs provided results of the degree of flexion, extension, rotation, and lateral flexion to the left. 270 participants were included in the assessment. Four studies were included for the evaluation of the ROM in flexion of which two studies showed significant differences. One favoring the MFR and one favoring the control group. All in all, there was no significant difference and a medium overall effect ( $d = 0.70$ , 95%CI = [-0.36 to 1.77],  $p = 0.15$ ). The in-between group comparison of extension based on four studies showed a medium overall effect and was rated as non-significant ( $d = 0.54$ , 95%CI = [-0.54 to 1.62],  $p = 0.24$ ). Similarities were found in four studies, that provided the outcomes of rotation to the left ( $d = 0.26$ , 95%CI = [-0.54 to 1.05],  $p = 0.42$ ) and lateral flexion to the left ( $d = 0.48$ , 95%CI = [-0.21 to 1.16],  $p = 0.12$ ). There was no identification of significant




Fig. 1. PRISMA- Flowchart of literature search.




Fig. 2. Results of a meta-analysis of VAS.

differences and a small overall effect size between the control and MFR group (see Fig. 4).

**Discussion**

This meta-analysis provides a comprehensive assessment of MFR efficacy in managing chronic neck pain, drawing evidence from ten RCTs. The results highlight a

significant reduction in pain, reflected in VAS scores. However, findings for PPT and ROM were inconclusive. In contrast to Guo et al.’s study (2023), which reported no significant differences in ROM, our meta-analysis identified slight significance, in lateral flexion and rotation to the right, though with slight variances (0.04 and 0.05, respectively) [39]. These marginal changes, especially the 0.03 minimal difference in VAS, may not be perceptible to patients, underscoring the importance for clinicians to




Fig. 3. Results of a meta-analysis of pressure pain threshold.

consider the minimal clinically important difference within the specific population under study. The observed significance in pain outcomes compared to the non-significant findings in PPT and ROM prompts careful consideration. Limited trials focusing on PPT may contribute to the absence of significant differences, raising questions about the clinical relevance of MFR for enhancing neck mobility, particularly given the heterogeneity in interventions. The incorporation of diverse interventions, such as a multimodal approach combining MFR with stability training, as demonstrated by previous studies, introduces clinical relevance but complicates the interpretation of results. The combined approach makes it challenging to isolate the specific effects of MFR alone [29,34]. Despite good-to-excellent methodological quality (see Table 1),

inconsistency between studies prompts a critical analysis of observed effects and their clinical significance. Treatment modality variations in control groups contribute to observed heterogeneity, necessitating more homogeneous study designs for enhanced outcome comparability. Additionally, the absence of blinding, except for Gauns & Gurudut (2018), introduces complexity, warranting careful consideration of potential biases, as seen in other systematic reviews [14].

Chronic pain is a pervasive healthcare problem in industrialized countries and is associated with high societal and individual costs [8,38]. Prolonged diagnostic processes remain a common burden, while conventional treatment modalities such as surgeries and medications, including opioids and analgesics, continue to be utilized




Fig. 4. Results of a meta-analysis of different range of motion outcomes.

despite limited evidence supporting their long-term efficacy and cost-effectiveness [8]. In contrast, MFR offers advantages regarding time efficiency and the absence of specialized equipment required, potentially contributing to cost reduction [31].

This study proposes that MFR may be an effective treatment for chronic neck pain. Nevertheless, definitive evidence supporting MFR treatments for chronic neck pain is lacking due to the existing shortage of studies, underscoring a significant scientific gap. Subsequent investigations should prioritize identifying independent predictors, conducting direct comparisons with established therapies, standardizing MFR protocols for enhanced reproducibility, and assessing long-term effects to foster a more comprehensive understanding of its efficacy.

### Funding Statement & Competing Interest

The research leading to these results has not received funding.

There was no declaration of interest in the form of financial competition or personal relationships, that could have appeared to influence the work reported in this paper.

### Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at [doi:10.1016/j.physio.2023.12.002](https://doi.org/10.1016/j.physio.2023.12.002).

### References

- [1] Fandim J, Nitzsche R, Michaleff Z, Oliviera Pena Costa L, Saragiotto B. The contemporary management of neck pain in adults. *Pain Manag* 2021;11(1):75–87.
- [2] Kazeminasab S, Nejadghaderi S, Amiri P, et al. Neck pain: global epidemiology, trends and risk factors. *BMC* 2022;23:26.
- [3] Treede R, Rief W, Barke A, et al. A classification of chronic pain for ICD-11. *Pain* 2015;156(6):1003–7.
- [4] Pelacios-Ceña D, Albaladejo-Vicente A, Hernández-Barrera V, et al. Female gender is associated with a higher prevalence of chronic neck pain, chronic low back pain, and migraine: results of the Spanish National Health Survey, 2017. *Pain Med* 2021;22(2):382–95.
- [5] Ezzati K, Ravarian B, Saberi A, Salari A, Reyhanian Z, Khakpour M, et al. Prevalence of cervical myofascial pain syndrome and its correlation with the severity of pain and disability in patients with chronic non-specific neck pain. *Arch Bone Jt Surg* 2021;9(2):230–4.
- [6] Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D. Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *EJP* 2006;10(4):287. –287.
- [7] Robert Koch Institut (RKI). Kopf- und Rückenschmerzen, Tumorschmerzen. *Chronische Schmerzen*. 2002;7:7–27.
- [8] Gatchel R, McGeary D, McGeary C, Lippe B. Interdisciplinary pain management. *Am Psychol* 2014;69(2):119–30.
- [9] Stecco A, Pirri C, Stecco C. Fascial entrapment neuropathy. *Clin Anat* 2019;32(1):883–90.
- [10] Bordoni B, Zanier E. Clinical and symptomatological reflections: the fascial system. *J Multidiscip Health* 2014;14(7):401–4011.
- [11] Davis C. Myofascial release. *Complementary Therapies in Rehabilitation: Evidence for Efficacy in Therapy, Prevention, and Wellness*. 3rd ed. New Jersey: SLACK; 2009.
- [12] Roland B, Decina P, Descarreaux M, et al. Evidence-based guidelines for the chiropractic treatment of adults with neck pain. *J Manip Physiol Ther* 2014;37(1):42–63.
- [13] McKenney K, Silclair A, Elder C, Hutchins A. Myofascial release as a treatment for orthopaedic conditions: a systematic review. *J Athl Train* 2013;48(8):522–7.
- [14] Ajimsha M, Al-Mudahka N, Al-Madzhar J. Effectiveness of Myofascial release techniques: Systematic review of randomized controlled trials. *J Bodyw Mov Ther* 2014;19(1):12–102.
- [15] Webb T, Rajendran D. Myofascial techniques: what are their effects on joint range of motion and pain? – a systematic review and meta-analysis of randomized controlled trials. *J Bodyw Mov Ther* 2016;20(3):682–99.
- [16] Wu Z, Wang Y, Xiangling Y. Myofascial release for chronic low back pain: a systematic review and meta-analysis. *Front Med (Lausanne)* 2021;28(8):697986.
- [17] Suarez-Rodriguez V, Fede C, Pirri C, et al. Fascial innervation: a systematic review of the literature. *Intern J Mol Sci* 2022;23(10):5674.
- [18] Page M, McKenzie J, Bossuyt P, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Syst Rev* 2021;10(1):89.
- [19] Cashin A, McAuley J. *Clinimetrics: Physiotherapy Evidence Database (PEDro) Scale*. *J Physiother* 2020;66(1):59.
- [20] Haefelin M, Elfering A. Pain assessment. *Eur Spine J* 2006;15(1):17–24.
- [21] Park G, Chan K, Bog S, Jung M, Ho J. Reliability and usefulness of the pressure pain threshold measurement in patients with myofascial pain. *Ann Rehabil Med* 2011;35(2):412–7.
- [22] Magee D. *Orthopaedic Physical Assessment*. 6th ed. Amsterdam: Elsevier Saunders; 2014.
- [23] Schwarzer G. Meta: An R package for meta-analysis. *R News*. 2007;7(3):40–45.
- [24] Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-test and ANOVAs. *Front Psychol* 2013;26(4):863.
- [25] Melsen W, Bootsma M, Rovers M, Bonten M. The effects of clinical and statistical heterogeneity on the predictive results from meta-analyses. *Clin Microbiol Infect* 2015;20(2):123–9.
- [26] Greenland S, Senn S, Rothman K, et al. Statistical tests, P values, confident intervals, and power: a guide to misinterpretations. *Eur J Epidemiol* 2016;31(4):337–50.
- [27] Röver C, Knapp G, Friede T. Hartung-Knapp-Sidik-Jonkman approach and its modification for random-effects meta-analysis with few studies. *BMC Med Res Method* 2015;14(15):99.
- [28] Namvar H, Olyaei G, Moghadam B, Hosseinfar M. Effect of myofascial release technique on pain, disability, maximum isometric contraction of the extensor muscles, and pressure pain threshold in patients with chronic nonspecific neck pain: double-blinded randomized clinical trial. *Int J Med Res Health Sci* 2016;5(7):500–6. 2016.
- [29] El-Gendy M, Lasheen Y, Rezkalla W. Multimodal approach of electrotherapy versus myofascial release in patients with chronic mechanical neck pain: a randomized controlled trial. *Physiother Q* 2019;27(4):6–12.
- [30] Rodríguez-Huguet M, Rodríguez-Almagro D, Rodríguez-Huguet P, Martín-Valero R, Lomas-Vega R. Treatment of neck pain with myofascial therapies: a single-blind randomized controlled trial. *J Manip Physiol Ther* 2020;43(2):160–70.
- [31] Rodríguez-Fuentes I, De Toro F, Rodríguez-Fuentes G, Machado de Oliveira I, Meijide-Fafde R, Fuentes-Boquete I. Is myofascial release therapy cost-effective when compared with manual therapy to treat workers' mechanical neck pain? *J Manip Physiol Ther* 2020;43(7):683–90.

- [32] Gauns S, Gurudut P. A randomized controlled trial to study the effect of gross myofascial release on mechanical neck pain referred to upper limb. *Int J Health Sci* 2018;12(5):51–9.
- [33] Bakar Y, Sertel M, Oztürk A, Tütün E, Tatarlı N, Ankaralı H. Short term effects of classic massage compared to connective tissue massage on pressure pain threshold and muscle relaxation response in women with chronic neck pain: a preliminary study. *J Manip Physiol Ther* 2014;37(6):415–21.
- [34] Celenay S, Kaya D, Akbayrak T. Cervical and scapulothoracic stabilization exercises with and without connective tissue massage for chronic mechanical neck pain: a prospective, randomized controlled trial. *Man Ther* 2015;21:144–50.
- [35] Rodríguez-Fuentes I, De Toro F, Rodríguez-Fuentes G, Machado de Oliveira I, Meijide-Fañde R, Fuentes-Boquete I. Myofascial release therapy in the treatment of occupational mechanical neck pain. *Am J Phys Med Rehabil* 2016;95(7):507–15.
- [36] Rodríguez-Huguet M, Gil-Salú J, Rodríguez-Huguet P, Cabrera-Afonso J, Lomas-Vega R. Effects of myofascial release on pressure pain threshold in patients with neck pain. *Am J Med Rehabil* 2018;97(1):16–22.
- [37] Retamal JJ, Seijo AF, Cintas JD, Rincón A, Bragado AC. Effects of instrumental, manipulative and soft tissue approaches for the sub-occipital region in subjects with chronic mechanical neck pain. A randomized controlled trial. *Int J Environ Res Public Health* 2021;18(16):8636.
- [38] Brück K, Jacobi K, Schmidt T. Fascial treatment versus manual therapy (HVLA) in patients with chronic neck pain: a randomized controlled trial. *J Back Musculoskelet Rehabil* 2021;34(6):997–1006.
- [39] Guo Y, Lv X, Zhou Y, Li Z, She H, Bai L, et al. Myofascial release for the treatment of pain and dysfunction in patients with chronic mechanical neck pain: Systematic review and meta-analysis of randomised controlled trials. *Clin Rehabil* 2023;37(4):478–93.

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

**ScienceDirect**

# QUESTIONNAIRE

D3(25)

## Effectiveness of myofascial release for adults with chronic neck pain: a meta-analysis

### INSTRUCTIONS

- Read through the article and answer the multiple-choice questions provided below.
- **Some questions may have more than one correct answer;** in which case you must mark **all** the correct answers.

### Introduction

**Question 1:** What is the estimated prevalence of chronic pain in Western societies?

- A: 5%
- B: 8%
- C: 12%
- D: 20%

**Question 2:** Which population is particularly affected by chronic neck pain?

- A: Young adults
- B: Middle-aged, office-bound males
- C: Middle-aged females with sedentary lifestyles
- D: Elderly individuals

**Question 3:** Which of the following statements are **TRUE**?

- A: An anatomical structure that is hypothesized to be strongly related to the intensity of pain is the fascial continuum
- B: Trauma, inflammations or infections, and structural imbalance of the body lead to a tightening of the fascial strain
- C: If this condition becomes long-term, the loss of physiological adaptability leads to a lack of flexibility, pain, and limitation in movement
- D: There is abundant scientific research available

**Question 4:** What are the primary goals of myofascial release (MFR)? To .....

- A: Increase blood flow to the neck
- B: Restore the optimal length of a muscle
- C: Alleviate pain
- D: Strengthen the neck muscles

**Question 5:** How is myofascial release (MFR) typically applied?

- A: Using a small foam roller
- B: Dry needling
- C: Applying low-load and long-duration stretch
- D: Using electrical stimulation

### Method

**Question 6:** Which muscles were examined for Pressure-Pain-Threshold (PPT) in the studies?

- A: Pectoralis
- B: Trapezius
- C: Sternocleidomastoid
- D: Deltoid

### Results

**Question 7:** What was the average duration of the treatment sessions in the studies on chronic neck pain?

- A: 10 minutes
- B: 23 minutes
- C: 34 minutes
- D: 45 minutes

**Question 8: TRUE or FALSE?**

"The total score for the rating of the risk of bias assessment and methodological quality of the ten RCTs ranged between seven to nine points."

- A: TRUE
- B: FALSE

**Question 9:** What was observed in the meta-analysis regarding pain reduction?

- A: No significant reduction in pain
- B: Slight increase in pain
- C: Significant reduction in pain
- D: Moderate reduction in pain

**Question 10:** What was the conclusion regarding the effectiveness of MFR on Range of Motion (ROM)?

- A: Highly effective
- B: No effect
- C: Significant differences
- D: Reduced ROM

## Discussion

**Question 11:** What was the main finding of the meta-analysis regarding MFR's efficacy in managing chronic neck pain?

- A: Significant reduction in pain
- B: Significant improvement in PPT
- C: Significant improvement in ROM
- D: No significant findings

**Question 12:** How did the meta-analysis findings on ROM compare to *Guo et al.*'s study?

- A: Both found significant improvements in ROM
- B: The meta-analysis found slight significance, while *Guo et al.* found no significant differences
- C: Both found no significant differences in ROM
- D: *Guo et al.* found significant improvements, while the meta-analysis did not

**Question 13:** Is the following statement **TRUE** or **FALSE**?

“Limited trials focusing on PPT may contribute to the absence of significant differences, raising questions about the clinical relevance of MFR for enhancing neck mobility, particularly given the heterogeneity in interventions”.

- A: TRUE
- B: FALSE

**Question 14:** What challenge does the incorporation of diverse interventions, such as combining MFR with stability training, introduce?

- A: Simplifies the interpretation of results
- B: Isolates the specific effects of MFR
- C: Complicates the interpretation of results
- D: Ensures homogeneity in study designs

**Question 15:** What does the study suggest about the current evidence supporting MFR treatments for chronic neck pain?

- A: There is definitive evidence supporting MFR
- B: There is a significant scientific gap due to a shortage of studies
- C: MFR is not effective for chronic neck pain
- D: MFR is more effective than conventional treatments

**THE END**



# FOCUS ON HEALTH

[www.foh-cpd.co.za](http://www.foh-cpd.co.za)

Cell: 074 230 3874

Tel: 012 653 0133 / 2373 / 2394

Mon-Fri: 08:00-16:00

400 Theuns van Niekerk Street

Wierda Park

0157

PO Box 71

Wierda Park

0149

## MEMBER INFO

(Complete sections marked with an asterisk \*)

**\*FOH number**

**\*HPCSA number**

**ID number**

**Personal detail changes**

**I hereby declare that the completion of this document is my own effort without any assistance.**

SIGNED:

DATE:

This activity is accredited for **THREE (3) CLINICAL CEU'S**. You have to achieve 70% to pass this activity.

## ANSWER SHEET

### D3(25)

Effectiveness of myofascial release for adults with chronic neck pain: a meta-analysis

	*Time spent on activity					hour					min					
	A	B	C	D	E	A	B	C	D	E	A	B	C	D	E	
1						8										
2						9										
3						10										
4						11										
5						12										
6						13										
7						14										
						15										

**How relevant was the content of this activity to your scope of practice?**

			
Not at all	Somewhat	Mostly	Extremely

### RETURN ANSWER SHEET TO

**Online members**, complete your answer sheet online

CPD PORTAL: <http://foh-cpd.co.za/cpd-online/cpd-user> (Username: FOH number & Password: ID)

(Results are available immediately)

**Email or hard copy members**, complete your answer sheet online or return it to:

**WHATSAPP:** 074 230 3874 **EMAIL:** info@foh-cpd.co.za

(Results will be sent via email)

You will receive a confirmation of receipt SMS within 12-24hours, if not received please send again.

Have a suggestion or comment?

Send it to us on **EMAIL** or **WHATSAPP**