



FOCUS ON  
**HEALTH**

## **Environmental Health**

### **Self-study CPD activity**

#### **Activity Reference Number**

**H9(25) 2028**

#### **Topic**

Red meat industry

#### **Article(s)**

Food safety incidents in the red meat industry: A review of foodborne disease outbreaks linked to the consumption of red meat and its products, 1991 to 2021

#### **Profession(s)**

Environmental Health Practitioner



Environmental Health Assistant

Food Inspector

***You can complete any one, or all the parts of this activity***

Part 1 is approved for **TWO (2) Clinical** Continuing Education Units (CEU's)

Part 2 is approved for **THREE(3) Clinical** Continuing Education Units (CEU's)



## Review

# Food safety incidents in the red meat industry: A review of foodborne disease outbreaks linked to the consumption of red meat and its products, 1991 to 2021

Dein Warmate<sup>\*</sup>, Bukola A. Onarinde

National Centre for Food Manufacturing, University of Lincoln, Holbeach PE12 7PT, UK



## ARTICLE INFO

## Keywords:

Pathogen  
Contributory factor  
Risk  
Strategy  
Management  
Abattoir  
Slaughter

## ABSTRACT

Red meat is a significant source of human nutrition, and the red meat industry contributes to the economy of nations. Nonetheless, there is a widespread global concern about public health issues posed by severe food safety incidents within the red meat industry. Most of these incidents are associated with foodborne disease outbreaks that impact individual consumers, food businesses and society. This study adopts a systematic search and review approach to identify three decades of published investigation reports of global foodborne disease outbreaks linked with the consumption of red meat and products made from them. The review aims to evaluate the critical features of these outbreak incidents to get insight into their contributing factors and root causes. In particular, this review discusses the transmission setting (origin of pathogenic agents), the food vehicles mostly incriminated, the causative pathogens (bacteria, viruses, and parasites) causing the most illnesses, and the most commonly reported contributing factors to the outbreaks. This information can help researchers and food business operators (FBOs) inform future risk assessment studies and support risk management activities in developing risk-mitigating strategies for the industry. Findings from this study suggest that implementing food safety management strategies which include adequate control measures at all stages of the food chain, from farm to fork, is imperative in preventing outbreak incidents. Of equal importance is the need for enhanced and sustained public education about the risk of foodborne illnesses associated with meat and its products whilst discouraging the consumption of raw meat products, especially by high-risk groups.

## 1. Introduction

### 1.1. Background

Food safety incidents within the red meat industry pose a severe threat to public health as they continue to be a widespread concern globally (Shang and Tonsor, 2017). These could be in the form of food contamination, which predominantly occurs during processing, distribution, retail, or catering (FSA, 2020), and could potentially lead to foodborne disease outbreak(s) from where these incidents are identified.

Food safety incidents could potentially lead to a food product being withdrawn or recalled and enforcement action taken at the implicated food establishment(s) following investigation and intelligence information from responsible authorities (Robertson et al., 2016). A single food safety incident, such as a foodborne disease outbreak, can have very grievous consequences, with losses of lives and businesses (Hussain

and Dawson, 2013). Most of the serious food safety issues resulting in consumer health problems and costly product withdrawals and recalls from the food supply chain of potentially contaminated meat products are associated with microbial hazards, especially bacterial pathogens (Sofos, 2008). Although various foods can serve as sources of foodborne illness, the literature suggests that meat and meat products are important sources of human infections (Nørrung et al., 2009). Several studies on outbreak investigations have reported red meat, such as beef, pork and products derived from them, as the culprit (Bélanger et al., 2015; Bryan, 1980; Jeffer et al., 2021; Omer et al., 2018).

For example, The US Department of Agriculture's Food Safety and Inspection Service (FSIS) investigated foodborne illness outbreaks associated with its regulated products – meat and poultry – from 2007 through 2012. From the investigation, 163 outbreaks were identified, associated with 4132 illnesses, 772 hospitalisations, and 19 deaths. Further, the report revealed that 105 outbreaks were linked to raw beef

<sup>\*</sup> Corresponding author.

E-mail address: [dwarmate@lincoln.ac.uk](mailto:dwarmate@lincoln.ac.uk) (D. Warmate).

<https://doi.org/10.1016/j.ijfoodmicro.2023.110240>

Received 23 July 2022; Received in revised form 12 April 2023; Accepted 28 April 2023

Available online 3 May 2023

0168-1605/© 2023 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

products sold commercially (Robertson et al., 2016). Similarly, the Centres for Disease Control and Prevention (CDC) reports that an estimated 48 million people get sick, 128,000 hospitalisations, and 3000 deaths occur every year nationwide in the USA from foodborne illnesses, most of which are related to the consumption of meat and meat products (CDC, 2011).

In 2012, Alberta Health Services, Canada, reported an *Escherichia coli* O157:H7 infection outbreak linked to the consumption of needle-tenderised beef steaks produced with beef from a meat plant in Alberta and purchased at a store in Edmonton, Alberta (Helmuth et al., 2019). Consequently, a staggering 4000 tons of beef products were recalled to control the outbreak, thus, making it the largest beef recall in Canadian history (Helmuth et al., 2019).

Another study reports that about 1060 people had become ill, and 216 died, between 2017 and 2018, in the largest recorded *Listeria monocytogenes* ST6 outbreak linked to the consumption of processed meat products from one of the largest food companies in South Africa - Tiger Brands. This food safety incident's impact was catastrophic, resulting in a significant reduction in consumer confidence in South Africa (Smith et al., 2019).

The aftermath of these incidents has a broader socio-economic impact beyond affected consumers. The financial costs to the food industry and state economy also can be substantial, particularly when product recalls, trade restrictions, litigations, and lost market share result from large-scale outbreaks (Limon et al., 2020). This fact was corroborated by another study that concludes that the estimated cost of food safety incidents for the economy of the United States is around \$7 billion per year derived from consumer notifications, removal of food from shelves, and payment of damages as a result of lawsuits (Hussain and Dawson, 2013).

It is noteworthy that not all food safety incidents result in disease outbreaks. However, the key is that there have to be concerns or risks about the food product's safety that may require intervention to protect consumers' interests. These concerns could be actual, potential, or perceived risks of illness associated with consuming food (FSANZ, 2021). For example, a meat slaughter and cutting establishment in the North of the UK was prosecuted and handed fines and costs totalling £266,000. The prosecution was occasioned by an incident in the plant where the Food Standard Agency (FSA) identified failings by the food business operator (FBO) to ensure the removal of specified parts of the animals, required by law, referred to as "Specified Risk Material (SRM)" (FSA, 2019).

In a similar event, another meat processing plant based in the Midlands in the UK was prosecuted and fined over £250,000. The food safety incident, in this case, involved the FBO removing meat that the FSA previously detained on suspicion of risk of contamination with the potential of the meat being injurious to health (FSA, 2019).

Given the severe consequences of food safety incidents associated with red meat and its impact on individuals, businesses and the wider society, it is imperative to get insight into the contributing factors and root causes of these incidents to aid in developing risk-mitigating strategies and countermeasures. However, this study focuses on foodborne disease outbreaks, although many diffuse foodborne diseases also exist. Thus, this study was conducted as a comprehensive review of published investigation reports and officially reported global foodborne disease outbreak incidents linked with the consumption of red meat and its products to evaluate the critical features and associated consequences of these outbreak incidents in the red meat industry.

The review highlights the food vehicles and related animal species mostly incriminated in the outbreaks, the foodborne pathogens (bacteria, viruses, and parasites) causing the most illnesses and identifies contributing factors to these outbreaks. Furthermore, this review strives to help researchers and FBOs, through this information, to inform future risk assessment studies and support risk management activities to prevent future outbreaks and protect public health and enhance food safety.

## 2. Materials and method

### 2.1. General approach

This study adopted a "systematic search and review" methodology to search peer-reviewed literature from four journal databases (Ebsco-Food Science Source, Google Scholar, PubMed, and Web of Science) to identify and summarise published reports of global outbreaks of foodborne disease linked to the consumption of red meat and their products, from 1991 to 2021. Two official websites, the Centers for Disease Control and Prevention and the European Centers for Disease Control and Prevention websites were reviewed. A total of 24,995 foodborne disease outbreaks related to different foods presented in summary reports from 1991 to 2020 were screened from these sites to identify 1628 outbreaks linked to red meat and its products. On the other hand, to extract data from the journal article databases, a review protocol was created outlining the search strategy, screening, selection criteria, and data extraction process.

#### 2.1.1. Search strategy

A six-step measure adopted from Rasmussen University Learning & Library Services (2018) was applied to achieve this:

**2.1.1.1. Extraction of the main ideas – identification of keywords.** Firstly, the search string was started by identifying the concepts and keywords central to the title or review statement: Food Safety Incidents, Red Meat, and Consumption were the key concepts of focus.

**2.1.1.1.1. Creation of a search menu.** Brainstorming a list of synonyms and related words to the key concepts was undertaken and subsequently added to the corresponding concepts row to create a search menu:

Concept 1: Food Safety Incident (Outbreak, foodborne disease, foodborne illness).

Concept 2: Red Meat (Beef; beef product, meat product, ham, burger, beef carcass, pork product, cattle, sheep, bovine).

Concept 3: Consumption (Intake; eat, ate, linked, associated).

Several search strings were created by linking concepts and synonyms with the Boolean logic operator. For example, (("outbreak") AND ("consumption" OR "link\*" OR "associated") AND ("meat" OR "beef" OR "ham" OR "goat" OR "sheep" OR "pork" OR "cow" OR "boar") NOT ("poultry" OR "fish" OR "vegetable")); (("foodborne disease") AND ("consumption" OR "link\*" OR "associated") AND ("meat" OR "beef" OR "ham" OR "goat" OR "sheep" OR "pork" OR "cow" OR "boar") NOT ("poultry" OR "fish" OR "vegetable")); (("outbreak") AND ("consumption" OR "link\*" OR "associated") AND ("meat" OR "beef" OR "ham" OR "goat" OR "sheep" OR "pork" OR "cow" OR "boar") NOT ("poultry" OR "fish" OR "vegetable")); (("foodborne illness\*") AND ("consumption" OR "link\*" OR "associated") AND ("meat" OR "beef" OR "ham" OR "goat" OR "sheep" OR "pork" OR "cow" OR "boar") NOT ("poultry" OR "fish" OR "vegetable"))).

The following filters/limiters were also applied to the databases:

- Full text included
- Abstract included
- Scholarly (peer-reviewed) journals
- Publication date: Between 1991 and 2021

The searches resulted in 349 hits for Ebsco-Food Science Source, 90 for Google Scholar, 618 for PubMed, and 775 for Web of Science. Journal articles retrieved by the search string were then exported from the databases and imported into Endnote X9 citation management software, where duplicate publications were identified and removed using the "find duplicates" function.

### 2.1.2. Relevance screening

After duplicate removal, the selected articles were exported from Endnote X9 citation management software and imported into Rayyan systematic review software to facilitate screening.

The titles were first reviewed in the screening process to check if selected articles were appropriate, and then all abstracts were screened for relevance. The initial screening of the title and abstract was based upon one key question: "Does the abstract investigate foodborne outbreak(s) associated with the consumption of red meat and products made from them?". Full texts of the relevant articles were further screened to satisfy the inclusion and exclusion criteria (Table 1).

### 2.1.3. Data extraction process

Data extraction was conducted on all selected articles that satisfied the predefined inclusion and exclusion criteria. The outbreak details were entered into a Microsoft Excel 2013 spreadsheet for review before data analysis. Data fields of interest extracted from the selected articles (data set) for further analysis include the following: The year the outbreak incident occurred; the country where it happened; the setting or place associated with the outbreak; the causative pathogen (aetiology); incriminated meat types and their products; the number of cases (people ill); the number of hospitalisations; deaths; cases/patients with Haemolytic Uraemic Syndromes (HUS); contributory factors.

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were followed in reporting the methods and results of this review (Fig. 1). However, the researchers did not conduct a meta-analysis as the study's focus was to summarise critical features of foodborne disease outbreaks. Meta-analysis is mainly conducted on randomised controlled trials (RCTs) or intervention studies (EFSA, 2010; Uetani et al., 2009).

## 3. Results

Table 2 presents key features of the review of journal databases and official websites. In contrast, Table 3 shows the descriptive summary and citation list of each investigated outbreak reported in selected journal articles from the databases. The data sources from the official sites were summary reports without details of specific incidents (CDC, 2022a; ECDC, 2022a, 2022b).

### 3.1. Transmission setting

Table 4 presents the classification of transmission settings associated with foodborne disease outbreaks reported in the scientific literature. Of the 101 outbreak incidents, most were linked to meat processing establishments, 42(42.4 %), followed by household settings, 25(24.9 %), which resulted in 67.4 % of the total outbreaks reported. The remaining

**Table 1**  
Inclusion and exclusion criteria.

| Inclusion criteria   | Exclusion criteria  |
|--|---|
| Language: English  | Language: Any language other than English.  |
| Product: The outbreak must be linked to the consumption of red meat and its products.  | Product: The outbreak was linked to the consumption of poultry, fish, vegetable (s) or products other than red meat and its products. |
| Study type: Full outbreak investigation with detailed information such as source, food vehicle, aetiology etc.<br>- Adequately described outbreaks of foodborne illnesses. | Study type: Summary studies that lacked sufficient details on individual outbreaks.   |
| Outcome: 2 or more confirmed cases of illness.   | Outcome: Single case of illness.  |

N/B: The outcome is necessary to satisfy the definition of an outbreak, i.e., an incident in which two or more persons experience a similar illness after exposure to a common source.

32.7 % of the outbreak incidents occurred in the other reported five categories of establishments.

### 3.2. Food vehicles

Table 5 presents the proportion of reported outbreaks, illnesses, hospitalisation, and deaths associated with the different meat types/species. Beef and pork were reported as the type/species of red meat most frequently associated with outbreaks, with beef (1077 or 62.35 %) coming first, followed by pork (555 or 32.1 %). This was followed by mixed species of meat (33 or 1.9 %).

Overall, the consumption of beef meat and its products resulted in more illnesses (23,907 or 57.7 %), hospitalisations (5324 or 49.8 %), and deaths (4792 or 47.6 %) than the consumption of pork meat and its products. The consumption of pork meat resulted in 14,977 or 36.1 % illnesses, 4886 or 45.7 % hospitalisations, and 4727 or 47.7 % deaths.

Nevertheless, the ratio beef:pork in number of outbreaks is around 2:1, and this lowers to around 1.5:1 for number of cases, and almost 1:1 for hospitalizations and deaths. Thus, pork appears to cause more cases and more severe disease within an average outbreak.

Table 6 presents a classification of the different food vehicles (products) derived from the different red meat types/species.

### 3.3. Etiologic agents (foodborne pathogens)

Shown in Table 7 is the proportion of reported outbreaks, illnesses, hospitalisations, and deaths associated with the eighteen foodborne pathogens identified. Most of the outbreaks were reported to be caused by *Salmonella* (469 or 27.1 %), followed by *E. coli* (414 or 23.9 %) and then *Clostridium* (294 or 17 %). In the same vein, *Salmonella* was associated with more illnesses or outbreak cases (13,469 or 32.5 %), hospitalisations (4349 or 40.7 %) and deaths (3826 or 38 %) than all the other pathogens. This was followed by *E. coli* and *Clostridium*.

Table 8 reveals that *Clostridium* is more associated with beef and its product (209 or 71.1 %) followed by pork (80 or 27.2 %). *E. coli* is also more associated with beef (380 or 91.9 %), followed by pork (12 or 27.2 %), and *Salmonella* is also more associated with beef (244 or 52.5 %).

### 3.4. Contributing factors

The factors that were contributory to the outbreaks are based on findings from the investigation reports from the scientific literature, as summary data from the official government websites did not give details of individual incidents or causes and hence were not used in the identification of contributory factors. Thus, the findings revealed that most of the outbreak incidents were associated with "Temperature abuse" 71/101(70.3 %), followed by contamination 49/101(48.5 %). The third was "cross-contamination", 40/101(39.6 %) but also, at least one contributory factor was associated with each foodborne disease outbreak.

## 4. Discussion

With a systematic search and review methodology, this study identified the contributing factors and critical features of foodborne disease outbreak incidents published in technical journals and official websites associated with the consumption of red meat and products made from them. Most of the elements have been identified in previous studies to cause foodborne disease outbreaks linked to other foods that are not red meat. However, to the best of the researchers' knowledge, this study is the first to undertake a comprehensive review of foodborne disease outbreaks associated with the consumption of red meat and its products from a global standpoint.

Available data revealed that the 1729 outbreaks identified over the 30-year period under review resulted in 41,438 illnesses (outbreak cases), 10,691 hospitalisations, and 10,063 deaths. This indicates that about 5 outbreaks, 115 illnesses, 30 hospitalisations and 28 deaths




Fig. 1. PRISMA flow chart detailing article screening and selection through the systematic search and review process.

**Table 2**  
Summary of key features of the outbreaks reported over three decades.

| Key features               | Database | Official site | Total  |
|----------------------------|----------|---------------|--------|
| Outbreaks                  | 101      | 1628          | 1729   |
| Illnesses (outbreak cases) | 6800     | 34,638        | 41,438 |
| Hospitalisations           | 909      | 9782          | 10,691 |
| Deaths                     | 254      | 9809          | 10,063 |
| Cases with HUS             | 91       |               |        |
| Minimum median age         | 2 yrs    |               |        |
| Maximum median age         | 89 yrs   |               |        |

occurred each month. These figures are of course indicative as it could have been more, probably due to underreporting. The reported median age of outbreak cases ranged from 2 yrs to 89 yrs, indicating that there is no age limit for contracting foodborne illness.

#### 4.1. Transmission setting or origin of infective agent (pathogens)

The place or setting where meat and its products were contaminated, pathogens survived and proliferated differed in each outbreak. These settings are categorised in Table 4. Reports highlight how meat and its product are either mishandled or mistreated in the cause of processing and or preparation, thus, resulting in an outbreak incident.

The findings of this study revealed that most of the reported outbreaks were linked to meat processing establishments followed by household settings. This reflects the potential risk posed by meat processing establishments. The peculiarity of the nature of operations within meat slaughter and processing establishments is no doubt a factor. Poor environmental hygiene, GMP, and bio-security measures within meat abattoirs are significant risk factors (Butt et al., 2021).

The most significant risk to meat safety within meat establishments is the presence of enteropathogenic bacteria capable of causing disease in the intestines or gastrointestinal tract of food-producing animals, which

**Table 3**  
Descriptive summary and citation list of each investigated outbreak report.

| Year | Location        | Setting                              | Etiologic agent               | Food incriminated               | Type of meat | Contributory factors   | No. of cases | Median age | Hospitalisation | Deaths | HUS cases | Recall | Reference                   |
|------|-----------------|--------------------------------------|-------------------------------|---------------------------------|--------------|--|--------------|------------|-----------------|--------|-----------|--------|-----------------------------|
| 1993 | Switzerland     | Food manufacturing establishment     | <i>Salmonella</i> Braenderup  | Meat pies                       | Beef         | Contamination; Poor GMP; Poor hygiene practices; Cross-contamination.            | 156          | 32         | 6               | 0      | NA.       | No     | Urfer et al. (2000)         |
| 1995 | Australia       | Church dinner                        | <i>Salmonella</i> Typhimurium | Roast pork meat                 | Pork         | Temperature abuse; Cross-contamination   | 22           | 22         | 1               | 0      | NA.       | No     | Delpech et al., 1998        |
| 1998 | Canada          | Church Supper                        | <i>Trichinella</i>            | Bear Burger                     | Bear         | Temperature abuse  | 8            | NA         | 0               | 0      | NA.       | No     | Nelson et al., 2003         |
| 1999 | USA             | Restaurant                           | <i>E. coli</i> O157:H7        | Beef Tacos                      | Beef         | Temperature abuse  | 13           | 12         | 5               | 0      | 3         | No     | Jay et al. (2004)           |
| 1999 | USA             | Meat processing establishment        | <i>L. monocytogenes</i>       | Fankfurters and deli meats      | Pork         | Contamination; Temperature abuse   | 108          | NA         | 0               | 14     | 0         | Yes    | Mead et al. (2006)          |
| 2001 | Denmark         | Household setting                    | <i>Trichinella</i>            | Walrus and Polar bear meat      | Bear         | Contamination; Temperature abuse   | 6            | NA         | 0               | 0      | 0         | No     | Møller et al. (2005)        |
| 2001 | Spain           | Meat slaughter/cutting establishment | <i>Trichinella</i>            | Domestic pig meat               | Pork         | Contamination; Temperature abuse   | 26           | NA         | 0               | 0      | 0         | No     | Cortés-Blanco et al. (2002) |
| 2001 | Slovakia        | Household setting                    | <i>Trichinella spiralis</i>   | Pork meat                       | Pork         | Contamination; Poor Hygiene practices; Temperature abuse                         | 23           | 34.4       | 6               | 0      | 0         | No     | Reiterova et al. (2007)     |
| 2001 | Germany         | Meat processing establishment        | <i>Salmonella</i> Muenchen    | Pork meat                       | Pork         | Contamination; Poor hygiene practices; Temperature abuse                         | 198          | NA         | NA              | NA     | NA        | No     | Buchholz et al. (2005)      |
| 2001 | Germany         | Meat processing establishment        | <i>Salmonella</i> Goldcoast   | Fermented Sausage               | Pork         | Poor GMP; Cross-contamination.   | 44           | 54         | NA.             | NA     | NA        | No     | Bremer et al. (2004)        |
| 2001 | USA             | Restaurant                           | <i>Salmonella</i> Uganda      | Roast pork meat                 | Pork         | Poor hygiene practices; Cross-contamination                                      | 24           | NA.        | NA              | NA     | NA        | No     | Jones et al. (2004)         |
| 2002 | Canada          | Household setting                    | <i>Trichinella</i>            | Bear meat                       | Bear         | Contamination; Temperature abuse   | 71           | 42         | 5               | 0      | 0         | No     | Schellenberg et al., 2003   |
| 2002 | USA             | Meat cutting establishment           | <i>E. coli</i> O157:H7        | Beef burger                     | Beef         | Contamination; Cross-contamination; Poor hygiene practices; Temperature control. | 9            | 14         | 3               | 0      | 3         | Yes    | Vogt and Dippold (2005)     |
| 2002 | Sweden          | Meat processing establishment        | <i>E. coli</i> O157:H7        | Fermented Sausage               | Beef         | Poor GMP; Temperature abuse  | 39           | 14         | NA              | NA     | 12        | No     | Sartz et al. (2008)         |
| 2002 | USA             | Meat processing establishment        | <i>E. coli</i> O157:H7        | Ground beef                     | Beef         | Contamination; Temperature abuse   | 18           | 15         | 7               | 0      | 5         | Yes    | CDC (2002)                  |
| 2003 | New Zealand     | Household setting                    | <i>Campylobacter jejuni</i>   | Pre-cooked cocktail sausages    | Pork         | Temperature abuse; Cross-contamination   | 3            | NA         | 0               | 0      | 0         | No     | Graham et al. (2005)        |
| 2003 | USA             | Meat processing establishment        | <i>Salmonella</i> Typhimurium | Ground beef                     | Beef         | Cross-contamination; temperature abuse   | 58           | 49         | 11              | 0      | 0         | No     | Dechet et al. (2006)        |
| 2003 | USA             | School potluck lunch                 | <i>Salmonella</i> Typhimurium | Ground beef                     | Beef         | Poor food hygiene practice; Cross-contamination; Temperature abuse.              | 47           | 8          | 2               | 0      | 0         | No     | McLaughlin et al. (2006)    |
| 2004 | Italy           | Household setting                    | <i>Salmonella</i> Typhimurium | Corallina Salami                | Pork         | Cross-contamination; Poor hygiene practice                                       | 63           | 7.5        | NA              | NA     | NA        | No     | Luzzi et al., 2007          |
| 2004 | Belgium         | Meat cutting establishment           | Hepatitis A Virus (HAV)       | Raw beef                        | Beef         | Cross-contamination; Poor hygiene practice; Temperature abuse.                   | 269          | 36.6       | 0               | 0      | 0         | No     | Robesyn et al. (2009)       |
| 2004 | Italy           | Household setting                    | <i>E. coli</i> O157           | Dry fermented Salami            | Pork         | Contamination  | 3            | 29         | 2               | 0      | 0         | No     | Conedera et al. (2007)      |
| 2004 | Italy           | Meat processing establishment        | <i>E. coli</i> O157           | Salami                          | Pork         | Contamination  | 2            | 60         | 2               | 0      | 0         | No     | Conedera et al. (2007)      |
| 2005 | The Netherlands | Mobile food caterers                 | <i>Salmonella</i> Typhimurium | Filet américain (Steak tartare) | Beef         | Temperature abuse  | 56           | NA         | 0               | 0      | NA.       | No     | Kivi et al. (2007)          |
| 2005 | France          | Food manufacturing establishment     | <i>E. coli</i> O157:H7        | Beef burger                     | Beef         | Poor hygiene practices; Cross-contamination; Temperature abuse.                  | 69           | 5          | 17              | 0      | 17        | Yes    | King et al. (2009)          |

(continued on next page)

Table 3 (continued)

| Year | Location                   | Setting                              | Etiologic agent                 | Food incriminated                            | Type of meat  | Contributory factors   | No. of cases | Median age | Hospitalisation | Deaths | HUS cases | Recall | Reference                  |
|------|----------------------------|--------------------------------------|---------------------------------|--|---------------|--|--------------|------------|-----------------|--------|-----------|--------|----------------------------|
| 2005 | France                     | Meat slaughter/cutting establishment | <i>Salmonella</i> Manhattan     | Cooked pork; minced beef                     | Mixed species | Contamination; Cross-contamination; Poor hygiene practices.                      | 69           | NA         | 3               | 0      | NA.       | No     | Noël et al. (2006)         |
| 2005 | The Netherlands            | Household setting                    | <i>E. coli</i> O157:H7          | Steak tartare                                | Beef          | Temperature abuse  | 33           | 24         | 7               | 0      | 0         | No     | Doorduyn et al. (2006)     |
| 2005 | Australia                  | Household setting                    | <i>Salmonella</i> Typhimurium   | Lamb Liver                                   | Lamb          | Cross-contamination; Temperature abuse   | 31           | 23         | 0               | 0      | 0         | No     | Hess et al. (2008)         |
| 2005 | Belgium                    | Meat slaughter establishment         | <i>Salmonella</i> Ohio          | Raw minced pork                              | Pork          | Poor GMP; Poor hygiene practices; Cross-contamination                            | 60           | 50         | 0               | 0      | 0         | No     | Bertrand et al. (2010)     |
| 2005 | USA                        | Meat processing establishment        | Norovirus                       | Delicatessen meat                            | Beef          | Poor hygiene practices; Poor GMP; Cross-contamination.                           | 57           | NA         | NA              | 0      | 0         | No     | Malek et al. (2009)        |
| 2005 | Denmark                    | Restaurant                           | <i>Salmonella</i> Typhimurium   | Carpaccio (thinly sliced raw fillet of beef) | Beef          | Contamination; Temperature abuse   | 32           | NA         | 11              | 0      | 0         | Yes    | Ethelberg et al. (2007)    |
| 2006 | Norway                     | Slaughter/cutting establishment      | <i>E. coli</i> O103             | Cured mutton sausage                         | Mutton        | Poor GMP; Cross-contamination; Poor hygiene; Contamination.                      | 17           | NA         | 14              | 1      | 10        | No     | Schimmer et al. (2008)     |
| 2006 | Norway                     | Household setting                    | <i>Yersinia enterocolitica</i>  | RTE-Brawn                                    | Pork          | Contamination  | 11           | 44         | 4               | 2      | 0         | No     | Grahek-Ogden et al. (2007) |
| 2006 | Germany                    | Meat processing establishment        | <i>L. monocytogenes</i>         | RTE-Scalded pork                             | Pork          | Cross-contamination; Poor hygiene practices.                                     | 16           | NA         | 2               | 0      | 0         | No     | Winter et al. (2009)       |
| 2007 | USA                        | Chain store butchery                 | <i>Salmonella</i> Newport       | Ground beef                                  | Beef          | Contamination; Cross-contamination; Poor hygiene practices; Temperature control. | 42           | 41         | 17              | 0      | 0         | No     | Schneider et al. (2011)    |
| 2007 | USA                        | Meat processing establishment        | <i>Salmonella</i> Typhimurium   | RTE Frozen Pot Pies                          | Beef          | Contamination; Temperature abuse   | 401          | 20         | 128             | 0      | 0         | Yes    | CDC (2008)                 |
| 2007 | Denmark                    | Meat processing establishment        | <i>E. coli</i> O26:H11          | Organic fermented beef sausage               | Beef          | Poor hygiene practices; Cross-contamination.                                     | 20           | 2          | 0               | 0      | 0         | Yes    | Ethelberg et al. (2009)    |
| 2008 | Denmark, Norway and Sweden | Meat slaughter/cutting establishment | <i>Salmonella</i> Typhimurium   | Danish pork meat meals, sausages etc.        | Pork          | Cross-contamination  | 37           | 54         | 0               | 4      | NA.       | No     | Bruun et al., 2009         |
| 2008 | Australia                  | Wine tavern                          | <i>L. monocytogenes</i>         | Jellied pork                                 | Pork          | Cross-contamination; temperature abuse   | 13           | 62         | 4               | 0      | 0         | No     | Pichler et al., 2009       |
| 2008 | The Netherlands            | Household setting                    | <i>E. coli</i> O157:H7          | Steak tartare                                | Beef          | Temperature abuse  | 20           | 41         | 7               | 0      | 0         | No     | Jager et al. (2009)        |
| 2008 | USA                        | Meat slaughter establishment         | <i>E. coli</i> O157 (STEC O157) | Ground beef                                  | Beef          | Contamination; Poor GMP; Poor hygiene practices; Temperature abuse               | 99           | 21         | 19              | 0      | 1         | Yes    | CDC (2010)                 |
| 2009 | Italy and Hungary          | Restaurant                           | <i>Salmonella</i> Goldcoast     | Salami Sandwich                              | Pork          | Cross-contamination  | 79           | 50         | 17              | 2      | NA.       | No     | Scavia et al., 2013        |
| 2009 | The Netherlands            | Household setting                    | <i>Salmonella</i> Typhimurium   | Steak tartare                                | Beef          | Temperature abuse  | 23           | 17         | 8               | 1      | 0         | No     | Whelan et al. (2010)       |
| 2009 | Denmark                    | Mobile food caterers                 | <i>L. monocytogenes</i>         | Beef meat                                    | Beef          | Temperature abuse  | 8            | 78         | 0               | 2      | 0         | No     | Smith et al. (2011)        |
| 2009 | France                     | Hotel                                | <i>Trichinella</i>              | Smoked warthog meat                          | Warthog       | Contamination; Temperature abuse   | 3            | NA         | 0               | 0      | 0         | No     | Dupouy-Camet et al. (2009) |
| 2010 | France                     | School Canteen                       | <i>Salmonella</i> Typhimurium   | Beef burger                                  | Beef          | Contamination; Temperature abuse   | 554          | NA         | 31              | 0      | 0         | Yes    | Raguenaud et al. (2012)    |
| 2010 | The Netherlands            | Household setting                    | <i>Salmonella</i> Typhimurium   | Ossenworst-Raw beef Sausage                  | Beef          | Temperature abuse  | 90           | NA         | 45              | 0      | 0         | No     | Friesema et al. (2012)     |
| 2010 | France                     | Meat processing establishment        | <i>Salmonella</i> Typhimurium   | Dried pork sausage                           | Pork          | Cross-contamination  | 110          | 89         | 20              | 0      | 0         | Yes    | Bone et al. (2010)         |
| 2010 | USA                        | School grounds                       | <i>E. coli</i> O157:H7          | Venison product                              | Deer          | Temperature abuse  | 29           | NA         | 2               | 0      | 0         | No     | Rounds et al. (2012)       |

(continued on next page)

Table 3 (continued)

| Year | Location    | Setting                               | Etiologic agent                | Food incriminated               | Type of meat  | Contributory factors  | No. of cases | Median age | Hospitalisation | Deaths | HUS cases | Recall | Reference                          |
|------|-------------|---------------------------------------|--------------------------------|---------------------------------|---------------|---|--------------|------------|-----------------|--------|-----------|--------|------------------------------------|
| 2011 | France      | Meat processing establishment         | <i>E. coli</i> O157            | Frozen beef burger              | Beef          | Poor hygiene practices; Cross-contamination; Temperature abuse.               | 18           | 3          | 18              | 0      | 18        | No     | King et al. (2014)                 |
| 2011 | USA         | Household setting                     | <i>Trichinella</i>             | Wild boar meat - Game           | Wild Boar     | Contamination; Temperature abuse; Poor food safety practice.                  | 6            | NA         | 1               | 0      | 0         | No     | Holzbauer et al. (2014)            |
| 2011 | Canada      | Catering Event                        | <i>E. coli</i> O157:H7         | Pork roast                      | Pork          | Contamination; Temperature abuse.   | 29           | 31         | 7               | 0      | 0         | No     | Trotz-Williams et al. (2012)       |
| 2011 | France      | Meat processing establishment         | <i>Salmonella Typhimurium</i>  | Pork sausage                    | Pork          | Poor GMP; Cross-contamination   | 337          | 10         | 0               | 0      | 0         | Yes    | Gossner et al. (2012)              |
| 2011 | India       | Community gathering                   | <i>Trichinella</i>             | Roast pork meat                 | Pork          | Temperature abuse   | 54           | NA         | 0               | 1      | 0         | No     | Sharma et al. (2014)               |
| 2011 | Denmark     | Retail chain (supermarket)            | <i>Salmonella Typhimurium</i>  | RTE meat-Smoked pork tenderloin | Pork          | Contamination   | 22           | NA.        | NA              | NA     | NA        | Yes    | Wójcik et al. (2012)               |
| 2012 | Norway      | Hotel                                 | <i>Clostridium perfringens</i> | Beef stew                       | Beef          | Temperature abuse   | 43           | 16         | 0               | 0      | NA.       | No     | Wahl et al., 2013                  |
| 2012 | England     | Food manufacturing establishment      | <i>L. monocytogenes</i>        | Pork pie                        | Pork          | Cross-contamination; Poor hygiene practices; Temperature control.             | 14           | NA         | 0               | 0      | 0         | Yes    | Awofisayo-okuyelu et al. (2016)    |
| 2012 | Vietnam     | Household setting                     | <i>Trichinella</i>             | Raw pork meat                   | Pork          | Contamination; Temperature abuse  | 24           |            | 6               | 0      | 0         | No     | Van de et al. (2012)               |
| 2012 | Canada      | Slaughter/ processing establishment   | <i>E. coli</i> O157:H7         | Ground beef                     | Beef          | Poor GMP; Cross-contamination; Contamination.                                 | 18           | 31         | 6               | 0      | 0         | Yes    | Currie et al. (2019)               |
| 2012 | New Zealand | Meat processing establishment         | <i>L. monocytogenes</i>        | RTE meat                        | Mixed species | Contamination   | 4            | NA         | NA              | 2      | NA        | No     | Rivas et al. (2019)                |
| 2012 | Serbia      | Funeral reception                     | <i>Trichinella</i>             | Smoked pork                     | Pork          | Temperature abuse   | 13           | NA         | 8               | 0      | 0         | No     | Popović-Dragonjić and Kocić (2018) |
| 2012 | Belgium     | Meat slaughter establishment          | <i>E. coli</i> O157:H7         | Steak tartare                   | Beef          | Contamination; Poor Hygiene practices; Temperature abuse                      | 24           | 31         | 15              | 0      | 5         | No     | Braeye et al. (2014)               |
| 2013 | Scotland    | Food manufacturing establishment      | <i>L. monocytogenes</i>        | Steak pie                       | Beef          | Cross-contamination; Poor hygiene; Bad manufacturing practices                | 3            | 87         | 3               | 0      | 0         | No     | Okpo et al., 2015                  |
| 2013 | USA         | Meat slaughter/ cutting establishment | <i>E. coli</i> O157:H7         | Beef burger                     | Beef          | Poor hygiene practices; Poor GMP; Cross-contamination; Temperature abuse.     | 24           | 25         | 8               | 0      | NA.       | No     | Torso et al. (2015)                |
| 2013 | France      | Wedding event                         | Hepatitis E Virus (HEV)        | Spit-Roasted Piglet             | Pork          | Contamination; Temperature abuse  | 17           | 45         | 2               | 0      | 0         | No     | Guillois et al. (2016)             |
| 2013 | Germany     | Meat processing establishment         | <i>Trichinella</i>             | Wild boar meat - Game           | Wild Boar     | Contamination; Poor GMP   | 21           | 52         | 0               | 0      | 0         | No     | Faber et al. (2015).               |
| 2013 | Germany     | Meat cutting establishment            | <i>Salmonella Typhimurium</i>  | Pork sausage, Minced pork       | Pork          | Contamination; Cross-contamination; Temperature abuse; Poor hygiene practices | 61           | 57         | 36              | 0      | 0         | No     | Alt et al. (2015)                  |
| 2013 | China       | Restaurant                            | <i>Clostridium botulinum</i>   | Smoked Pork ribs                | Pork          | Contamination; Temperature abuse; Poor food safety practices.                 | 12           | NA         | 7               | 0      | 0         | Yes    | Feng et al. (2015)                 |
| 2013 | Germany     | Slaughter establishment               | <i>Salmonella</i> Infantis     | Raw minced pork                 | Pork          | Contamination; Poor GMP; Poor hygiene practices; Temperature abuse            | 267          | 56         | 85              | 0      | 0         | No     | Schroeder et al. (2016)            |
| 2014 | France      | Meat processing establishment         | <i>Salmonella Enteritidis</i>  | frozen beef burger              | Beef          | Contamination; Temperature abuse  | 45           | 9          | 0               | 0      | NA.       | Yes    | Jones et al. (2016)                |
| 2014 | Belgium     | Restaurant                            | <i>Trichinella</i>             | Wild boar meat                  | Wild Boar     | Contamination; Temperature abuse  | 16           | 37         | 10              | 0      | 0         | Yes    | Messiaen et al. (2016)             |

(continued on next page)

Table 3 (continued)

| Year | Location        | Setting                              | Etiologic agent                    | Food incriminated                        | Type of meat  | Contributory factors   | No. of cases | Median age | Hospitalisation | Deaths | HUS cases | Recall | Reference                                   |
|------|-----------------|--------------------------------------|------------------------------------|--|---------------|--|--------------|------------|-----------------|--------|-----------|--------|---|
| 2014 | Germany         | Pig Farm                             | <i>Salmonella</i> Muenchen         | Raw pork products, sausages and spreads. | Pork          | Poor hygiene practices   | 247          | 56         | 0               | 0      | 0         | No     | <a href="#">Schielke et al. (2017)</a>      |
| 2014 | Sweden          | Meat processing establishment        | <i>L. monocytogenes</i>            | Cold cut ham                             | Pork          | Cross-contamination; Poor hygiene practices  | 51           | 77         | 0               | 0      | 0         | Yes    | <a href="#">Dahl et al. (2017)</a>          |
| 2014 | Germany         | Meat processing establishment        | <i>L. monocytogenes</i>            | RTE Sausage                              | Pork          | Poor GMP; Poor hygiene practices; Cross-contamination                                      | 39           | 73         | 25              | 18     | 0         | Yes    | <a href="#">Lachmann et al. (2021)</a>      |
| 2015 | Italy           | Food manufacturing establishment     | <i>L. monocytogenes</i>            | Hog head cheese (Processed pork)         | Pork          | Cross-contamination; Poor hygiene practice   | 35           | 75         | 24              | 4      | 0         | Yes    | <a href="#">Duranti et al., 2018</a>        |
| 2015 | Spain           | Household setting                    | Hepatitis E Virus (HEV)            | Wild boar meat                           | Wild Boar     | Contamination; Temperature abuse   | 8            | NA         | 0               | 0      | 0         | No     | <a href="#">Rivero-Juarez et al. (2017)</a> |
| 2015 | France          | Meat slaughter/cutting establishment | <i>Trichinella</i>                 | Figatelli (pork sausages)                | Pork          | Contamination; Cross-contamination; Poor hygiene practices.                                | 3            | NA         | 1               | 0      | 0         | No     | <a href="#">Ruetsch et al. (2016)</a>       |
| 2015 | Scotland        | Slaughter/cutting establishment      | <i>E. coli</i> O157:H7             | Venison                                  | Deer          | Cross-contamination; Temperature abuse   | 12           | 41         | 0               | 0      | 0         | No     | <a href="#">Smith-Palmer et al. (2018)</a>  |
| 2015 | England         | Meat processing establishment        | <i>E. coli</i> O157                | RTE meat                                 | Mixed species | Cross-contamination; Poor hygiene practices; Temperature control.                          | 15           | 38         | 10              | 0      | 7         | No     | <a href="#">Wilson et al. (2018)</a>        |
| 2015 | USA             | Meat slaughter/cutting establishment | <i>Salmonella</i> Infantis         | Roaster pork meat                        | Pork          | Contamination; Cross-contamination; Poor GMP; Poor hygiene practices; Temperature control. | 192          | 35         | 30              | 0      | 0         | Yes    | <a href="#">Kawakami et al. (2019)</a>      |
| 2015 | The Netherlands | Meat processing establishment        | <i>Salmonella Typhimurium</i>      | Filet américain (raw beef spread)        | Beef          | Contamination; Temperature abuse   | 45           | 21         | 29              | 0      | 0         | No     | <a href="#">Freidl et al. (2018)</a>        |
| 2016 | Italy           | School Canteen                       | <i>L. monocytogenes</i>            | Cooked beef ham                          | Beef          | Cross-contamination; Temperature abuse   | 162          | NA         | 5               | 0      | 0         | No     | <a href="#">Maurella et al. (2018)</a>      |
| 2016 | The Netherlands | Meat processing establishment        | <i>Salmonella Bovismorbificans</i> | Ham                                      | Pork          | Contamination  | 54           | 65         | 15              | 0      | 0         | Yes    | <a href="#">Brandwagt et al. (2018)</a>     |
| 2016 | Japan           | Restaurant                           | <i>Trichinella T9</i>              | Bear meat meal                           | Bear          | Contamination; Temperature abuse; Poor food safety practice.                               | 21           | 19         | 0               | 0      | 0         | No     | <a href="#">Tada et al. (2018)</a>          |
| 2016 | Serbia          | Household setting                    | <i>Trichinella britovi</i>         | Wild boar Ham and sausages               | Wild Boar     | Contamination; Temperature abuse   | 114          | NA         | 19              | 0      | 0         | No     | <a href="#">Dmitric et al. (2018)</a>       |
| 2016 | Italy           | Household setting                    | <i>Trichinella</i>                 | Boar meat sausage                        | Wild Boar     | Contamination; Temperature abuse   | 5            | NA         | 1               | 0      | 0         | No     | <a href="#">Turiac et al. (2017)</a>        |
| 2016 | Switzerland     | Meat processing establishment        | <i>L. monocytogenes</i>            | Meat Pate                                | Beef          | Cross-contamination; Poor GMP; Poor hygiene practices; Temperature abuse                   | 7            | 84.8       | 0               | 0      | 0         | Yes    | <a href="#">Althaus et al. (2017)</a>       |
| 2016 | India           | Household setting                    | <i>Aeromonas hydrophila</i>        | Carcase meat                             | Beef          | Contamination; Temperature abuse   | 33           | NA.        | NA              | NA     | NA        | No     | <a href="#">Tsheten et al. (2016)</a>       |
| 2016 | Japan           | Household setting                    | <i>E. coli</i> O157:H7             | Raw minced meat cutlets                  | Mixed species | Temperature abuse  | 61           | 31         | 24              | 0      | 4         | Yes    | <a href="#">Furukawa et al. (2018)</a>      |
| 2016 | Canada          | Household setting                    | <i>Trichinella</i>                 | Jerky (dried bear meat)                  | Bear          | Temperature abuse  | 10           | NA         | 0               | 0      | 0         | No     | <a href="#">Dalcin et al. (2017)</a>        |
| 2017 | England         | Household setting                    | <i>E. coli</i> O157:H7             | Frozen beef burger                       | Beef          | Temperature abuse  | 12           | 16         | 8               | 0      | 4         | Yes    | <a href="#">Byrne et al. (2020)</a>         |
| 2017 | Uganda          | Meat slaughter/cutting establishment | <i>Anthrax</i>                     | Beef meat                                | Beef          | Poor GMP; Poor hygiene practices; Contamination; Temperature abuse                         | 61           | NA         | 0               | 0      | 0         | No     | <a href="#">Nakanwagi et al. (2020)</a>     |
| 2017 | South Africa    | Meat processing establishment        | <i>L monocytogenes</i>             | RTE meat - Polony sausage                | Mixed species | Poor GMP; Poor hygiene practices; Contamination; Cross-contamination.                      | 937          | NA         | 2               | 197    | NA        | Yes    | <a href="#">Thomas et al. (2020)</a>        |

(continued on next page)

Table 3 (continued)

| Year | Location        | Setting                              | Etiologic agent               | Food incriminated                           | Type of meat  | Contributory factors   | No. of cases | Median age | Hospitalisation | Deaths | HUS cases | Recall | Reference                                |
|------|-----------------|--------------------------------------|-------------------------------|---|---------------|--|--------------|------------|-----------------|--------|-----------|--------|--|
| 2017 | USA             | Retreat                              | <i>Toxoplasma gondii</i>      | Venison - grilled                           | Deer          | Temperature abuse  | 11           | 51         | 0               | 0      | 0         | No     | <a href="#">Schumacher et al. (2021)</a> |
| 2017 | USA             | House party                          | <i>Trichinella</i>            | Raw boar meat                               | Wild boar     | Temperature abuse  | 36           | NA         | 0               | 0      | 0         | No     | <a href="#">Heaton et al. (2018)</a>     |
| 2018 | Denmark         | Household setting                    | <i>Salmonella Typhimurium</i> | Medister sausage (Raw Danish pork sausage). | Pork          | Cross-contamination; temperature abuse                             | 49           | 66         | 0               | 0      | 0         | No     | <a href="#">Helmuth et al., 2019</a>     |
| 2018 | USA             | Meat slaughter/cutting establishment | <i>Salmonella Newport</i>     | Beef meat                                   | Beef          | Contamination; Poor GMP; Poor hygiene practices; Temperature abuse | 255          | 36         | 60              | 2      | 0         | No     | <a href="#">Plumb et al. (2019)</a>      |
| 2018 | Canada          | Hunting Retreat                      | <i>Toxoplasma gondii</i>      | Venison                                     | Deer          | Temperature abuse  | 6            | NA         | 1               | 0      | 0         | No     | <a href="#">Gaulin et al. (2020)</a>     |
| 2019 | The Netherlands | Food manufacturing establishment     | <i>L. monocytogenes</i>       | RTE meat products                           | Mixed species | Cross-contamination; Poor hygiene practices.                       | 21           | 78         | 21              | 3      | 0         | Yes    | <a href="#">ECDC-EFSA (2019)</a>         |
| 2019 | China           | Household setting                    | <i>Clostridium botulinum</i>  | Vacuum-packed Ham                           | Pork          | Temperature abuse  | 4            | 55         | 4               | 1      | 0         | No     | <a href="#">Min et al. (2021)</a>        |
| 2019 | The Netherlands | Household setting                    | <i>E. coli O157</i>           | Steak tartare                               | Beef          | Temperature abuse  | 20           | 41         | 7               | 0      | 0         | No     | <a href="#">Greenland et al. (2009)</a>  |
| 2019 | UK.             | Retail chain (supermarket)           | <i>E. coli O157:H7</i>        | Raw minced beef                             | Beef          | Contamination; Temperature abuse                                   | 13           | 19         | 5               | 0      | 2         | No     | <a href="#">Butt et al. (2021)</a>       |

**Table 4**

Classification of reported transmission settings associated with foodborne disease outbreak.

| Transmission setting  | Category                         |
|---|----------------------------------|
| Meat cutting plant, pig farm, red meat abattoir, meat slaughterhouse, fresh meat processing establishment, game meat cutting plant, slaughter/co-located cutting plant, slaughter establishment, retail chain butcher, privately owned meat butchery, local meat butcher store. | Meat processing establishment    |
| Food manufacturing establishment  | Food manufacturing establishment |
| Church supper, church dinner, school canteen, school grounds, school potluck lunch  | Institutional establishment      |
| Hotel, restaurant, mobile food caterers, hospital catering  | Foodservice establishment        |
| Chain store butchery, retail chain (supermarket)  | Retail chain supermarket         |
| Home, house, house party  | Household setting (HHS)          |
| Wedding events; catering events; wine tavern; hunting retreats; funeral receptions; community gatherings; family retreats.  | Others (OTR)                     |

**Table 5**

The proportion of outbreaks, illnesses, hospitalisations, and deaths associated with the different meat-types.

| Meat species  | Outbreak    | Illness       | Hosp.         | Death         |
|---------------|-------------|---------------|---------------|---------------|
|               | No.         | No.           | No.           | No.           |
| Bear          | 21          | 202           | 30            | 25            |
| Beef          | 1077        | 23,907        | 5324          | 4792          |
| Bison         | 3           | 76            | 49            | 31            |
| Buffalo       | 2           | 15            | 3             | 3             |
| Deer          | 4           | 58            | 3             | 0             |
| Goat          | 6           | 85            | 27            | 27            |
| Lamb          | 17          | 261           | 116           | 118           |
| Mixed species | 33          | 1620          | 197           | 328           |
| Mutton        | 1           | 17            | 14            | 1             |
| Pork          | 555         | 14,977        | 4886          | 4727          |
| Warthog       | 1           | 3             | 0             | 0             |
| Wild Boar     | 9           | 217           | 42            | 11            |
| <b>Total</b>  | <b>1729</b> | <b>41,438</b> | <b>10,691</b> | <b>10,063</b> |

**Table 6**

Classification of incriminated food vehicles associated with foodborne disease outbreak incidents.

| Incriminated food (meat products) vehicle   | Category                              |
|---|---------------------------------------|
| Bear Burger, Bear meat, Beef burger, Beef meat, Boar meat sausage, Carcase meat, Carpaccio (thinly sliced raw fillet of beef), Domestic pig meat, Filet américain (raw beef spread), Ground beef, Lamb Liver, Medister sausage (Raw Danish pork sausage), minced beef, Minced pork, Ossenworst-Raw beef Sausage, Pork meat, Pork sausage, Raw beef, Raw boar meat. Raw minced beef, Raw minced pork, Raw pork meat, Roaster pork meat, sausages and spreads, Steak tartare, Venison, Walrus and Polar bear meat, Wild boar Ham and sausages, Wild boar meat | Freshly processed meat products (FPM) |
| Corallina Salami, cured mutton sausage, Delicatessen meat, Dried pork sausage, Dry fermented Salami, Fermented Sausage, Figatelli (pork sausages), Frankfurters and deli meats, Jerky (dried bear meat) Meat Pate, Organic fermented beef sausage, RTE-Brawn, RTE-meat products, RTE-Polony sausage, RTE-Scalded pork, RTE-Smoked pork tenderloin, Salami Sandwich, Smoked pork   | Ready-to-eat meat products (RTEM)     |
| Beef stew, Beef Tacos, Cold cut ham, cooked beef ham, Frozen Pot Pies, Grilled venison, Hog head cheese (processed Pork), Jellied pork, Meat pies, Pork pie, Pork roast, Pre-cooked cocktail sausages, Roast pork meat, Smoked pork ribs, smoked warthog meat, Spit-Roasted Piglet, Steak pie, vacuum-packed ham  | Cooked meat/meal products (CMM)       |

**Table 7**

The proportion of outbreaks, illnesses, hospitalisation, and deaths caused by etiologic agents.

| etiologic agent               | Outbreak    | Illnesses     | Hospitals     | Deaths        |
|-------------------------------|-------------|---------------|---------------|---------------|
|                               | No          | No            | No            | No            |
| <i>Aeromonas hydrophila</i>   | 1           | 33            | 0             | 0             |
| <i>Anthrax</i>                | 1           | 61            | 0             | 0             |
| <i>Bacillus cereus</i>        | 51          | 692           | 279           | 279           |
| <i>Campylobacter jejuni</i>   | 15          | 261           | 44            | 42            |
| <i>Clostridium</i>            | 294         | 11,071        | 2762          | 2843          |
| <i>E. coli</i>                | 414         | 4953          | 1092          | 791           |
| <i>Enterobacter</i>           | 2           | 30            | 0             | 0             |
| Hepatitis E Virus             | 1           | 269           | 0             | 0             |
| Hepatitis E Virus             | 2           | 25            | 2             | 240           |
| <i>Listeria monocytogenes</i> | 120         | 3106          | 627           | 530           |
| Norovirus                     | 144         | 3067          | 429           | 442           |
| <i>Salmonella</i>             | 469         | 13,469        | 4349          | 3826          |
| <i>Sapovirus</i>              | 1           | 22            | 22            | 22            |
| <i>Shigella</i>               | 7           | 173           | 0             | 0             |
| <i>Staphylococcus</i>         | 134         | 3048          | 652           | 662           |
| <i>Toxoplasma gondii</i>      | 2           | 17            | 1             | 0             |
| <i>Trichenella</i>            | 55          | 953           | 426           | 382           |
| <i>Yersinia</i>               | 16          | 188           | 6             | 4             |
| <b>Total</b>                  | <b>1729</b> | <b>41,438</b> | <b>10,691</b> | <b>10,063</b> |

renders them symptomless carriers of foodborne diseases among live-stock presented for slaughter (Lianou et al., 2017). For instance, some meat-borne pathogens (such as *Salmonella*, *Campylobacter*, *E. coli* O157:H7, and *Yersinia*) live in the intestine of animals that frequently and easily contaminate meat (DeWaal et al., 2006). Some of these pathogens are found on the skins of animals, potentially causing cross-contamination of meat and the environment (Antic et al., 2021). In addition to the skin, the gastrointestinal and respiratory tracts, urine, and milk duct are other vital sources of infection (Nel et al., 2004).

Contamination of meat can also occur during processing at abattoirs, particularly during evisceration, where gut contents are likely to contaminate exposed meat if incorrect techniques are applied (noted from the researcher's professional experience). However, the literature reveals that carcass contamination during beef slaughter and processing mostly occurs during dehiding and skinning (Antic et al., 2021; Kennedy et al., 2014; McEvoy et al., 2000) and preventing cross-contamination from the hide surface to the carcass is very challenging due to the nature of the hide removal process (McEvoy et al., 2000). Other potential sources of contamination during the dressing of meat carcasses include the knives, hands and clothing of workers, processing equipment such as saws, boning tables, conveyors, and water used to wash carcasses (Govender et al., 2013). Therefore, it is vital to observe a minimum standard of sanitation and personal hygiene to minimise contamination.

#### 4.2. Food vehicles most incriminated in the outbreaks

For clarity and simplicity, the various types of meats and their products implicated as vehicles of foodborne disease were classified into three broad categories (Table 6). These include freshly processed meat, ready-to-eat meat and cooked meat/meal products (CMM).

Freshly processed meat products are primarily raw meat that requires further processing, in most cases, before consumption. For instance, some freshly processed meat products, such as steak tartar and beef burger, were consumed raw or undercooked, i.e., cooked "medium-rare" (Kivi et al., 2007).

The different ways these foods are processed or prepared can affect how they are contaminated or influence the survival and proliferation of pathogens (Bryan et al., 1997).

Regarding the species or type of meat, the data reveals that beef (1077 or 62.35 %) and pork (555 or 32.1 %) were most frequently associated with outbreaks (Table 5). Thus, beef and pork meat and their products were the most incriminated food vehicles reported during the 30-year period under review. This is partly connected to beef and pork

**Table 8**  
Proportion of etiologic agent associated with meat type/species.

| Etiologic agent               | Bear | Beef | Bison | Buffalo | Deer | Goat | Lamb | Mixed | Mutton | Pork | Warthog | Wild Boar |
|-------------------------------|------|------|-------|---------|------|------|------|-------|--------|------|---------|-----------|
|                               | No   | No   | No    | No      | No   | No   | No   | No    | No     | No   | No      | No        |
| <i>Aeromonas hydrophila</i>   | 0    | 1    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 0    | 0       | 0         |
| <i>Anthrax</i>                | 0    | 1    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 0    | 0       | 0         |
| <i>Bacillus cereus</i>        | 0    | 36   | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 4    | 0       | 0         |
| <i>Campylobacter jejuni</i>   | 0    | 8    | 0     | 0       | 0    | 0    | 0    | 15    | 0      | 6    | 0       | 0         |
| <i>Clostridium</i>            | 0    | 209  | 0     | 0       | 0    | 1    | 0    | 4     | 0      | 80   | 0       | 0         |
| <i>E. coli</i>                | 0    | 380  | 3     | 1       | 2    | 1    | 9    | 5     | 1      | 12   | 0       | 0         |
| <i>Enterobacter</i>           | 0    | 2    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 0    | 0       | 0         |
| Hepatitis A Virus             | 1    | 0    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 0    | 0       | 0         |
| Hepatitis E Virus             | 0    | 0    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 1    | 0       | 1         |
| <i>Listeria monocytogenes</i> | 0    | 62   | 0     | 0       | 0    | 0    | 0    | 6     | 0      | 50   | 0       | 0         |
| Norovirus                     | 0    | 86   | 0     | 0       | 0    | 0    | 0    | 3     | 0      | 58   | 0       | 0         |
| <i>Salmonella</i>             | 0    | 244  | 0     | 0       | 0    | 1    | 8    | 6     | 0      | 207  | 0       | 0         |
| <i>Sapovirus</i>              | 0    | 0    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 1    | 0       | 0         |
| <i>Shigella</i>               | 0    | 7    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 6    | 0       | 0         |
| <i>Staphylococcus</i>         | 0    | 40   | 0     | 0       | 0    | 0    | 0    | 4     | 0      | 91   | 0       | 0         |
| <i>Toxoplasma gondii</i>      | 0    | 0    | 0     | 0       | 2    | 0    | 0    | 0     | 0      | 0    | 0       | 0         |
| <i>Trichinella</i>            | 18   | 0    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 21   | 1       | 8         |
| <i>Yersinia</i>               | 0    | 0    | 0     | 0       | 0    | 0    | 0    | 0     | 0      | 15   | 0       | 0         |

being the most consumed species of red meat globally. According to the FAO (2019), pork is the most widely eaten meat in the world (36 %), followed by poultry (33 %), beef (24 %), and goats/sheep (5 %).

The consumption of beef meat and its products showed to have resulted in more illnesses or outbreak cases (23,907 or 57.7 %), and hospitalisations (5324 or 49.8 %) than the consumption of pork meat and its products.

#### 4.3. Foodborne pathogens (bacteria, viruses, and parasites) causing the most illnesses

Of the eighteen foodborne pathogens identified, *Salmonella*, *E. coli*, and *Clostridium* were reported as the three most common causes of foodborne outbreaks linked to the consumption of red meat and its products, respectively (Table 7).

##### 4.3.1. Etiologic agent vs meat type

Findings in this study regarding etiologic agents associated with meat types/species reveal that specific pathogens are more associated with specific meat types and their products (Table 8). For instance, it was revealed that *Clostridium* was more associated with beef and its product (71.1 %), followed by pork (27.2 %); *E. coli* was also more associated with beef (91.9 %) followed by pork (12 or 27.2 %) and *Salmonella* was also more associated with beef (52.45 %) followed by pork (44.4 %). This knowledge can inform risk management decisions. More so, specific interventions directed at these food categories and their associated pathogens would help prevent or reduce the frequency of foodborne illness outbreaks.

##### 4.3.2. *Salmonella*

In this analysis, *Salmonella* caused most (469 or 27.1 %) of the outbreaks, with consequent illnesses or outbreak cases of 13,469 or 32.5 %, 4349 or 40.7 % hospitalisations and 3826 or 38 % deaths. In many countries, food poisoning outbreaks caused by *Salmonella* in meat and its products are commonly reported, and *Salmonella* is considered one of the foodborne pathogens of great concern (Plumb et al., 2019). A previous study on an overview of foodborne outbreaks in Canada for 16 years highlighted the importance of *Salmonella* as an etiologic agent of enteric illness as it was the most common pathogen and caused 40.3 % of the outbreaks (Bélanger et al., 2015).

*Salmonella* is a bacterium that can cause an illness or a zoonotic disease called salmonellosis in humans. In the EU, over 91,000 salmonellosis cases are reported each year (EFSA, 2022). Food-source animals sometimes become infected, or their skin or feet become contaminated

on farms. The salmonellae are conveyed to processing plants in or on these animals, usually in faecal material on their feet. These organisms can spread to carcasses or cuts of meat during processing. In most cases, when raw or inadequately cooked, these products allow salmonellae to survive and cause foodborne disease outbreaks (Bryan, 1980).

Salmonellosis may result in severe illness; however, the spread of the disease can be controlled by hygienic slaughtering practices and thorough cooking and refrigeration of meat and its products (Braeye et al., 2014).

##### 4.3.3. *E. coli*

The second-highest number of outbreaks (414), was attributed to *E. coli*, resulting in 4953 illnesses, 1092 hospitalisations and 791 deaths. Like *Salmonella*, *E. coli* is a naturally occurring bacteria commonly found in the gastrointestinal tract of animals, especially cattle (Wahl et al., 2013; Brusa et al., 2022). Most *E. coli* strains are harmless, although Shiga toxin-producing *E. coli* (STEC), the strain most commonly reported in this study, can cause severe and life-threatening foodborne diseases such as haemolytic uraemic syndrome (HUS), thrombocytopenia, renal failure, and occasionally death, especially in children and the elderly (Bélanger et al., 2015; WHO, 2022).

This study shows that 100 % of all the outbreak cases with HUS reported from the scientific literature reviewed were associated with *E. coli* infection, which corroborates relevant literature. For instance, a study conducted by Feitz et al. (2021) reveals that Shiga toxin-producing *E. coli* bacteria infection was implicated to have caused 90 % of childhood haemolytic uraemic syndrome (HUS) cases.

Most of the outbreak investigation reports suggest that *E. coli* infection transmission to consumers occurs through the consumption of raw or undercooked meat (such as minced or ground beef burgers) that had already been contaminated during slaughter and processing in meat processing plants (Duranti et al., 2018). Faecal contamination of carcass meat and cross-contamination from carcasses to the hands of operators, equipment, tools and the abattoir environment and from them to other carcasses are also risk factors (Costa et al., 2020; Dickson and Acuff, 2017). Thus, good manufacturing/hygiene practices in meat processing establishments should help prevent beef product contamination (Buncic et al., 2014). As STEC is heat-sensitive and destroyed by thorough cooking, meat preparation at home should adhere to basic food safety practices such as “cook thoroughly”.

##### 4.3.4. *Clostridium*

*Clostridium* was the third most reported pathogen incriminated in this study, following *E. coli*, however the findings reveal it caused more

illnesses (11071), hospitalisations (2762) and deaths (2843) than *E. coli*. *Clostridium perfringens* are spore-forming bacteria that can survive heat, dryness, and other environmental conditions. *C. perfringens* bacteria are one of the most common causes of food poisoning, with an estimated nearly 1 million foodborne illnesses caused by the bacterium every year in the United States (CDC, 2022b).

Chicken, turkey and red meat (such as beef and pork) are specific foods commonly associated with *C. perfringens* outbreaks. This is especially when they are cooked in large batches and held at unsafe temperatures, which occurs predominantly in settings like schools, hospitals, factories and catering establishments (Bennet et al., 2013).

On the other hand, *Clostridium botulinum* is a bacterium that causes botulism. The bacterium produces heat-resistant endospores which are very resistant to a number of environmental stresses, such as heat and high acid (CDC, 2013). Although the bacteria and spores alone do not cause disease, their production of botulinum toxin renders them pathogenic (Feng et al., 2015).

Foodborne botulism results from the ingestion of preformed botulinum toxin in food. The toxin can be found in food that has not been properly cooked, processed, handled, or canned and is often present in canned food, such as vegetables, meat, and seafood products (CDC, 2013).

#### 4.4. Most reported contributing factors to the outbreaks

Several factors contributed to meat contamination and the survival or proliferation of the etiologic agents in the meat and its products. These factors contributed to the outbreaks' causation and were identified and classified as: contamination, cross-contamination, poor "GMP", poor hygiene practice, temperature abuse, and poor food safety practice. At least one contributory factor was associated with each foodborne disease outbreak. The most reported contributing factor was "Temperature abuse". This result suggests that temperature control is a critical factor in preventing outbreaks.

The findings indicate that it may take more than one contributing factor to cause an outbreak and that all contributing factors may not occur in one establishment. For instance, contamination might have happened in a different setting (e.g., meat processing establishment, food service establishment, retail chain store, etc.), and mishandling that allowed survival or proliferation (e.g., temperature abuse; cross-contamination) might have occurred in another setting in an outbreak incident.

Some of the reported factors that have contributed to outbreaks classed as "temperature abuse" include improper cooling, insufficient temperature or time during cooking, improper hot holding and insufficient temperature or time during reheating of food.

Besides the temperature abuse, contamination was reported at different stages of the production and preparation of meat and its products. Poor standards of hygiene (environment and personal), poor GMP and cross-contamination in meat processing establishments have contributed to and influenced the occurrence of foodborne outbreak incidents. Others were mishandling or mistreatment of meat coupled with a disregard for suitable hygienic measures and a lack of food safety awareness and training of food handlers. These factors contributed to pathogens coming into contact with food and, in some cases, enabling the survival and multiplication of pathogens to cause foodborne disease outbreak incidents.

#### 5. Strengths and limitations

A key strength of this study is its contributions and novelty in undertaking a comprehensive review of factors associated with the consumption of red meat and its products from a global standpoint. This type of review combines the strengths of a critical review with an exhaustive systematic search process which can provide a much more complete picture of the prevalence of the research problem in question

(Grant and Booth, 2009). The search was conducted following an organised plan or stipulated set of clearly defined objectives and reproducible methods. Literature reveals several types of systematic reviews that adhere to various guidelines on conducting a review to help standardise them and improve their quality (Kitchenham, 2004; Needleman, 2002).

One crucial aspect that cuts across these guidelines is, assessing the methodological quality of the included studies or considering the validity of the individual studies (Ahn and Kang, 2018; EFSA, 2010; Kitchenham, 2004; Needleman, 2002). However, this review did not undertake a quality assessment of selected outbreak reports as part of the inclusion/exclusion criteria partly because this review does not adhere to a specific systematic review guideline.

Nevertheless, the researchers retrieved the outbreak reports included in this study from peer-reviewed literature, which implies that the journal articles have gone through an evaluation process in which journal editors and other expert scholars critically assessed the quality and validity of the research methodology and procedures. The Peer-review process ensures that the research quality and the findings' validity are high. Besides, the review of official websites ensures data credibility given that government websites provide accurate records of foodborne disease outbreaks.

However, the findings of this study have some limitations. First, reporting and publication bias may have resulted in under-reporting foodborne disease outbreaks in the literature, given that large outbreaks are more likely to be investigated and published. Since the terminology used to describe outbreaks report investigations differed by report, it is possible that outbreak investigation reports that did not explicitly mention the keywords used in the search strings could have been missed in the database searches. However, the robust search strategy adopted for this study was aimed to minimise the number of potentially missed outbreak reports.

Another consideration is that because the search strategy, by design, did not include searches beyond bibliographic databases, such as exploring the grey literature, it might have missed some vital outbreak reports in the grey literature. Further, as English language was one of the inclusion criteria in this study, publications in other languages other than English language might have been missed as well.

#### 6. Conclusion

With a systematic search and transparent review methodology, this review identifies outbreak investigation reports of global foodborne disease associated with red meat consumption for three decades, from 1991 through 2021. The study provided a comprehensive account of the features of the reported outbreaks under review. Eighteen pathogens (of which *Salmonella* was the most reported) were identified from these outbreaks to have caused 1729 outbreaks, 41,438 illnesses, 10,691 hospitalisations, and 10,063 deaths. Besides, these outbreak incidents were reported to have led to product recalls/withdrawals, which are costly to food establishments due to downtime. Beyond affected consumers, these outbreak incidents negatively impact the socio-economy of businesses and countries. It is noteworthy that many of the foodborne diseases reported are diffuse cases. For instance, outbreaks generated due to contamination originating at meat processing establishments have been reported to result in widespread contamination through the supply chain and affected a potentially much greater number of people across geographically distributed locations with a massive public health impact.

Given that a combination of factors can contribute to the same outbreak, preventing outbreak incidents requires approaches or control measures at all stages of the food chain. Thus, implementing robust food safety management strategies from agricultural production on the farm to processing, manufacturing, and preparing foods in all establishments - commercial, institutional and household establishments - is imperative.

Findings from this study have important implications for improving

food safety programmes, setting priorities based on preventive intervention for future outbreaks, and strengthening food safety policies, including risk assessment, especially in the red meat industry. Besides, the data can inform decisions at places and operations where regulations need to be established and enforced and indicate educational and training needs.

Suffice it to say that since the findings indicate meat processing establishments are chiefly implicated as the transmission setting or origin of pathogens, abattoirs have a significant role to play in measures toward the reduction of outbreak incidents in general. However, a shared responsibility of all stakeholders within the food chain: from farm to fork, i.e., farmers, processors, distributors, retailers, caterers, and consumers, all have their part to minimise the risks of food safety incidents.

In a nutshell, to prevent outbreak incidents and tackle food safety challenges faced by the meat industry, the findings of this review highlight the need for a high meat-producing standard which includes maintaining adequate hygiene standards and process controls during the slaughter and processing of animals. The need for adequate information, training and supervision of meat production/handling operatives in safe food handling practices to prevent cross-contamination and proliferation of foodborne pathogens is of equal importance.

The findings also point to the need for enhanced and sustained public education about the risk of foodborne illnesses associated with meat and its products whilst discouraging the consumption of raw meat products, especially by high-risk groups.

## Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## CRediT authorship contribution statement

DW and BO: Conceptualisation of the overarching research goals and aims.

DW and BO: Design of the data search method.

DW: Data Collation and data presentation.

DW and BO: Data analysis and data interpretation.

DW: Preparation and writing of the initial draft.

DW and BO: Structuring of manuscript and critically reviewing the contents.

BO: Supervision; oversight and leadership responsibility for the research activity planning and execution.

BO: Approval to submit the manuscript for publication.

All authors have read and agreed to the published version of the manuscript.

## Declaration of competing interest

The authors declare no conflict of interest.

## Data availability

Data will be made available on request.

## Acknowledgments

We thank Prof. Mark Swainson and Dr. Samson Oyeyinka for editing and proofreading the final manuscript draft.

## References

- Ahn, E., Kang, H., 2018. Introduction to systematic review and meta-analysis. *Korean J. Anesthesiol.* 71 (2), 103.
- Alt, K., Simon, S., Helmeke, C., Kohlstock, C., Prager, R., Tietze, E., Rabsch, W., Karagiannis, I., Werber, D., Frank, C., Fruth, A., 2015. Outbreak of uncommon O4

- non-agglutinating salmonella typhimurium linked to minced pork, Saxony-Anhalt, Germany, January to April 2013. *Plos One* 10 (6), e0128349.
- Althaus, D., Jermini, M., Giannini, P., Martinetti, G., Reinholz, D., Nüesch-Inderbinen, M., Lehner, A., Stephan, R., 2017. Local outbreak of listeria monocytogenes serotype 4b sequence type 6 due to contaminated meat pâté. *Foodborne Pathog. Dis.* 14 (4), 219–222.
- Antic, D., Houf, K., Michalopoulou, E., Blagojevic, B., 2021. Beef abattoir interventions in a risk-based meat safety assurance system. *Meat Sci.* 182, 108622.
- Awofisayo-okuyelu, A., Arunachalam, N., Dallman, T., Grant, A., Aird, H., McLauchlin, J., Painsset, A., Amar, C., 2016. An outbreak of human listeriosis in England between 2010 and 2012 associated with the consumption of pork pie. *J. Food Prot.* 79 (5), 732–740.
- Bélanger, P., Tanguay, F., Hamel, M., Phypers, M., 2015. Foodborne illness: an overview of foodborne outbreaks in Canada reported through outbreak summaries: 2008–2014. *Can. Commun. Dis. Rep.* 41 (11), 254.
- Bennet, S.D., Walsh, K.A., Gould, L.H., 2013. Foodborne disease outbreaks caused by *Bacillus cereus*, *Clostridium perfringens*, and *Staphylococcus aureus*—United States, 1998–2008. *Clin. Infect. Dis.* 57 (3), 425–433.
- Bertrand, S., Dierick, K., Heylen, K., De Baere, T., Pochet, B., Robesyn, E., Lokietek, S., Van Meerven, E., Imberechts, H., De Zutter, L., Collard, J.M., 2010. Lessons learned from the management of a national outbreak of salmonella Ohio linked to pork meat processing and distribution. *J. Food Prot.* 73 (3), 529–534.
- Bone, A., Noel, H., Le Hello, S., Pihier, N., Danan, C., Raguenaud, M.E., Salah, S., Bellali, H., Vaillant, V., Weill, F.X., Jourdan-da Silva, N., 2010. Nationwide outbreak of *Salmonella enterica* serotype 4, 12: ii-infections in France, linked to dried pork sausage, March–May 2010. *Euro Surveill.* 15 (24), 19592.
- Braeye, T., Denayer, S., De Rauw, K., Forier, A., Verluuyten, J., Fourie, L., Dierick, K., Botteldoorn, N., Quoilin, S., Cosse, P., Noyen, J., 2014. Lessons learned from a textbook outbreak: EHEC-O157: H7 infections associated with the consumption of raw meat products, June 2012, Limburg, Belgium. *Arch Public Health* 72 (1), 1–7.
- Brandwagt, D., van den Wijngaard, C., Tulen, A.D., Mulder, A.C., Hofhuis, A., Jacobs, R., Heck, M., Verbruggen, A., van den Kerkhof, H., Slegers-Fitz-James, I., Mughini-Gras, L., 2018. Outbreak of *Salmonella Bovismorbificans* associated with the consumption of uncooked ham products, the Netherlands, 2016 to 2017. *Euro Surveill.* 23 (1), 17–00335.
- Bremer, V., Leitmeyer, K., Jensen, E., Metzger, U., Meczulat, H., Weise, E., Werber, D., Tschaep, H., Krienbrock, L., Glaser, S., Ammon, A., 2004. Outbreak of salmonella goldcoast infections linked to consumption of fermented sausage, Germany 2001. *Epidemiol. Infect.* 132 (5), 881–887.
- Brusa, V., Restovich, V., Galli, L., Arias, R., Linares, L., Costa, M., Díaz, V.R., Pugin, D., Leotta, G., 2022. Reduction of Shiga toxin-producing *Escherichia coli* in a beef abattoir. *Food Sci. Technol. Int.* 28 (1), 50–59.
- Bruun, T., Sørensen, G., Forshell, L.P., Jensen, T., Nygård, K., Kapperud, G., Lindstedt, B. A., Berglund, T., Wingstrand, A., Petersen, R.F., Müller, L., Kjølseth, C., Ivarsson, S., Hjertqvist, M., Löfdahl, S., Ethelberg, S., 2009. An outbreak of *Salmonella typhimurium* infections in Denmark, Norway and Sweden, 2008. *Euro Surveill.* 14 (10), 9147.
- Bryan, F.L., 1980. Foodborne diseases in the United States associated with meat and poultry. *J. Food Prot.* 43 (2), 140–150.
- Bryan, F.L., Guzewich, J.J., Todd, E.C., 1997. Surveillance of foodborne disease III. Summary and presentation of data on vehicles and contributory factors; their value and limitations. *J. Food Prot.* 60 (6), 701–714.
- Buchholz, U., Brodhun, B., Brockmann, S.O., Dreweck, C.M., Prager, R., Tschäpe, H., Ammon, A., 2005. An outbreak of salmonella München in Germany associated with raw pork meat. *J. Food Prot.* 68 (2), 273–276.
- Buncic, S., Nychas, G.J., Lee, M.R., Koutsoumanis, K., Hebraud, M., Desvaux, M., et al., 2014. Microbial pathogen control in the beef chain: recent research advances. *Meat Sci.* 97 (3), 288–297.
- Butt, S., Smith-Palmer, A., Shand, A., McDonald, E., Allison, L., Maund, J., Fernandes, A., Vishram, B., Greig, D.R., Jenkins, C., Elson, R., 2021. Evidence of on-going transmission of Shiga toxin-producing *Escherichia coli* O157: H7 following a foodborne outbreak. *Epidemiol. Infect.* 1–25.
- Byrne, L., Kaindama, L., Bentley, M., Jenkins, C., Aird, H., Oliver, I., Paranthaman, K., 2020. Investigation into a national outbreak of STEC O157: H7 associated with frozen beef burgers, UK, 2017. *Epidemiol. Infect.* 148.
- Centres for Disease Control and Prevention (CDC), 2002. Multistate outbreak of *Escherichia coli* O157: H7 infections associated with eating ground beef—United States, June–July 2002. *Morb. Mortal. Wkly Rep.* 51 (29), 637–639.
- Centres for Disease Control and Prevention (CDC), 2008. Multistate outbreak of salmonella infections associated with frozen pot pies—United States, 2007. *Morb. Mortal. Wkly Rep.* 57 (47), 1277–1280.
- Centres for Disease Control and Prevention (CDC), 2010. Two multistate outbreaks of Shiga toxin-producing *Escherichia coli* infections linked to beef from a single slaughter facility—United States, 2008. *Morb. Mortal. Wkly Rep.* 59 (18), 57–560.
- Centres for Disease Control and Prevention (CDC), 2011. Foodborne Burden. Available from: <https://www.cdc.gov/foodborneburden/burden>.
- Centers for Disease Control and Prevention (CDC), 2013. National enteric disease surveillance: botulism surveillance overview. Available from: <https://www.cdc.gov/botulism/surveillance.html>.
- Centres for Disease Control and Prevention (CDC), 2022a. Prevent Illness From *C. Perfringens*. Available from: <https://www.cdc.gov/foodsafety/diseases/clostridium-perfringens.html>.
- Centres for Disease Control and Prevention (CDC), 2022a. National outbreak reporting systems. Available from: <https://www.cdc.gov/nors/index.html>.
- Conedera, G., Mattiazzi, E., Russo, F., Chiesa, E., Scorzato, I., Grandesso, S., Bessegato, A., Fioravanti, A., Caprioli, A., 2007. A family outbreak of *Escherichia*

- coli O157 haemorrhagic colitis caused by pork meat salami. *Epidemiol. Infect.* 135 (2), 311–314.
- Cortés-Blanco, M., García-Cabañas, A., Guerra-Peguero, F., Ramos-Aceitero, J.-M., Herrera-Guibert, D., Martínez Navarro, J.F., 2002. Outbreak of trichinellosis in Cáceres, Spain, December 2001–February 2002. *Euro Surveill.* 7 (10), 2–5.
- Costa, M., Pracca, G., Sucari, A., Galli, L., Ibarogoyen, J., Gentiluomo, J., et al., 2020. Comprehensive evaluation and implementation of improvement actions in bovine abattoirs to reduce pathogens exposure. *Prev. Vet. Med.* 176, 104933.
- Currie, A., Honish, L., Cutler, J., Locas, A., Lavoie, M.C., Gaulin, C., Galanis, E., Tschetter, L., Chui, L., Taylor, M., Jamieson, F., 2019. Outbreak of *Escherichia coli* O157: H7 infections linked to mechanically tenderised beef and the largest beef recall in Canada, 2012. *J. Food Prot.* 82 (9), 1532–1538.
- Dahl, V., Sundqvist, L., Hedenström, I., Löfdahl, M., Alm, E., Ringberg, H., Lindblad, M., Wallensten, A., Thisted Lambert, S., Jernberg, C., 2017. A nationwide outbreak of listeriosis associated with cold-cuts, Sweden 2013–2014. *Infect. Ecol. Epidemiol.* 7 (1), 1324232.
- Dalcin, D., Zarlenga, D.S., Larter, N.C., Hoberg, E., Boucher, D.A., Merrifield, S., Lau, R., Ralevski, F., Cheema, K., Schwartz, K.L., Boggild, A.K., 2017. *Trichinella nativa* outbreak with rare thrombotic complications associated with meat from a black bear hunted in northern Ontario. *Clin. Infect. Dis.* 64 (10), 1367–1373.
- Dechet, A.M., Scallan, E., Gensheimer, K., Hoekstra, R., Gunderman-King, J., Lockett, J., Wrigley, D., Chege, W., Sobel, J., Multistate Working Group, 2006. Outbreak of multidrug-resistant *Salmonella enterica* serotype Typhimurium definitive type 104 infection linked to commercial ground beef, northeastern United States, 2003–2004. *Clin. Infect. Dis.* 42 (6), 747–752.
- Delpach, V., McNulty, J., Morgan, K., 1998. A salmonellosis outbreak linked to internally contaminated pork meat. *Aust. N. Z. J. Public Health* 22 (2), 243–246.
- DeWaal, C.S., Hicks, G., Barlow, K., Alderton, L., Vegosen, L., 2006. Foods associated with foodborne illness outbreaks from 1990 through 2003. *Food Prot. Trends* 26 (7), 466–473.
- Dickson, J.S., Acuff, G.R., 2017. Maintaining the safety and quality of beef carcass meat. In: Acuff, G.R., Dickson, J.S. (Eds.), *Ensuring Safety and Quality in the Production of Beef*. Burleigh Dodds Science Publishing, Cambridge, pp. 145–167.
- Dmitric, M., Debeljak, Z., Vidanovic, D., Sekler, M., Vaskovic, N., Matovic, K., Karabasil, N., 2018. *Trichinella britovi* in game meat linked to human trichinellosis outbreak in Serbia. *J. Parasitol.* 104 (5), 557–559.
- Doorduyn, Y., De Jager, C.M., Van Der Zwaluw, W.K., Friesema, I.H., Heuvelink, A.E., De Boer, E., Wannet, W.J.B., van Duynhoven, Y.T.H.P., 2006. Shiga toxin-producing *Escherichia coli* (STEC) O157 outbreak, the Netherlands, September–October 2005. *Euro Surveill.* 11 (7), 5–6.
- Dupouy-Camet, J., Lecam, S., Talabani, H., Ancelle, T., 2009. Trichinellosis acquired in Senegal from warthog ham, march 2009. *Euro Surveill.* 14 (21), 19220.
- Durant, A., Sabbatucci, M., Blasi, G., Acciari, V.A., Ancora, M., Bella, A., Busani, L., Centorame, P., Cammà, C., Conti, F., De Medici, D., 2018. A severe outbreak of listeriosis in Central Italy with a rare pulsotype associated with processed pork products. *J. Med. Microbiol.* 67 (9), 1351–1360.
- ECDC, EFSA (European Centre for Disease Prevention and Control, European Food Safety Authority), 2019. Multi-country outbreak of *Listeria monocytogenes* sequence type 6 infections linked to ready-to-eat meat products – 25 November 2019. Available from: <https://www.ecdc.europa.eu/en/publications-data/rapid-outbreak-assessment-multi-country-outbreak-listeria-monocytogenes-sequence>.
- EFSA (European Food Safety Authority), 2010. Guidance for those carrying out systematic reviews. Application of systematic review methodology to food and feed safety assessments to support decision making. *EFSA J.* 8 (6), 1637.
- Ethelberg, S., Smith, B., Torpdahl, M., Lisby, M., Boel, J., Jensen, T., Nielsen, E.M., Mølbak, K., 2009. Outbreak of non-O157 Shiga toxin-producing *Escherichia coli* infection from consumption of beef sausage. *Clin. Infect. Dis.* 48 (8), 78–81.
- Ethelberg, S., Sørensen, G., Kristensen, B., Christensen, K., Krusell, L., Hempel-Jørgensen, A., Perge, A., Nielsen, E.M., 2007. Outbreak with multi-resistant *salmonella typhimurium* DT104 linked to carpaccio, Denmark, 2005. *Epidemiol. Infect.* 135 (6), 900–907.
- European Centre for Disease Prevention and Control (ECDC), 2022. Trichinellosis. Available from: <https://www.ecdc.europa.eu/en/trichinellosis/facts>.
- European Food Safety Authority (EFSA), 2022. Zoonotic Disease: Salmonella (Fact Sheet). Available from: <https://www.efsa.europa.eu/en/topics/topic/salmonella>.
- European Centre for Disease Prevention and Control (ECDC), 2022. Monitoring and Reporting Data and Trends. Available from: <https://ecdc.europa.eu/en/publications-data>.
- Faber, M., Schink, S., Mayer-Scholl, A., Ziesch, C., Schönfelder, R., Wichmann-Schauer, H., Stark, K., Nöckler, K., 2015. Outbreak of trichinellosis due to wild boar meat and evaluation of the effectiveness of post exposure prophylaxis, Germany, 2013. *Clin. Infect. Dis.* 60 (12), 98–104.
- Feitz, W.J., Bouwmeester, R., van der Velden, T.J., Goorden, S., Licht, C., van den Heuvel, L.P., van de Kar, N.C., 2021. The Shiga toxin receptor globotriaosylceramide as a therapeutic target in Shiga toxin E. coli mediated HUS. *Microorganisms* 9 (10), 2157.
- Feng, L., Chen, X., Liu, S., Zhou, Z., Yang, R., 2015. Two-family outbreak of botulism associated with the consumption of smoked ribs in Sichuan Province China. *Int. J. Infect. Dis.* 30, 74–77.
- Freidl, G., Schoss, S., Te Wierik, M., Heck, M., Tolsma, P., Urbanus, A., Slegers-Fitz-James, I., Friesema, I., 2018. Tracing back the source of an outbreak of *Salmonella typhimurium*; national outbreak linked to the consumption of raw and undercooked beef products, the Netherlands, October to December 2015. *PLoS Curr.* 10 (2018) <https://doi.org/10.1371/2Fcurrents.outbreaks.1c667d62b51eb9840f57eb617e56bc1>.
- Friesema, I.H., Schimmer, B., Ros, J.A., Ober, H.J., Heck, M.E., Swaan, C.M., de Jager, C.M., Peran i Sala, R.M., van Pelt, W., 2012. A regional *Salmonella enterica* serovar Typhimurium outbreak associated with raw beef products, The Netherlands, 2010. *Foodborne Pathog. Dis.* 9 (2), 102–107.
- FAO (Food and Agriculture Organisation of the United Nations), 2019. Market and trade commodities/intergovernmental group on meat and dairy products <https://www.fao.org/markets-and-trade/commodities/meat/intergovernmental-group-on-meat-and-dairy-products/en/>. <https://steakholderfoods.com/what-are-the-most-consumed-meats-in-the-world/>.
- FSA (Food Standard Agency), 2019 & 2020. Food incidents, product withdrawals and recalls. Last updated 31 December 2020. <https://www.food.gov.uk/business-guidance/food-incidents-product-withdrawals-and-recalls>.
- FSANZ (Food Standards Australia & New Zealand), 2021. Food Incidents. <https://www.foodstandards.gov.au/industry/foodrecalls/Pages/Food-Incidents.aspx>.
- Furukawa, I., Suzuki, M., Masaoka, T., Nakajima, N., Mitani, E., Tasaka, M., Teranishi, H., Matsumoto, Y., Koizumi, M., Ogawa, A., Oota, Y., 2018. An outbreak of enterohemorrhagic *Escherichia coli* O157: H7 infection associated with minced meat outlets in Kanagawa, Japan. *Jpn. J. Infect. Dis.* 71 (6), 436–441.
- Gaulin, C., Ramsay, D., Thivierge, K., Tataryn, J., Courville, A., Martin, C., Cunningham, P., Désilets, J., Morin, D., Dion, R., 2020. Acute toxoplasmosis among Canadian deer hunters associated with consumption of undercooked deer meat hunted in the United States. *Emerg. Infect. Dis.* 26 (2), 199.
- Gossner, C.M., Van Cauteren, D., Le Hello, S., Weill, F.X., Terrien, E., Tessier, S., Janin, C., Brisabois, A., Dusch, V., Vaillant, V., Jourdan-da Silva, N., 2012. Nationwide outbreak of *salmonella enterica* serotype 4, [5], 12: i-infection associated with consumption of dried pork sausage, France, November to December 2011. *Euro Surveill.* 17 (5), 20071.
- Govender, R., Naidoo, D., Buys, E.M., 2013. Managing meat safety at south african abattoirs. *Int. J. Agric. Biol. Sci. Eng.* 7, 3843.
- Graham, C., Whyte, R., Gilpin, B., Cornelius, A., Hudson, J.A., Morrison, D., Graham, H., Nicol, C., 2005. Outbreak of campylobacteriosis following pre-cooked sausage consumption. *Aust. N. Z. J. Public Health* 29 (6), 507–510.
- Grahek-Ogden, D., Schimmer, B., Cudjoe, K.S., Nygård, K., Kapperud, G., 2007. Outbreak of yersinia enterocolitica serogroup O:9 infection and processed pork Norway. *Emerg. Infect. Dis.* 13 (5), 754.
- Grant, M.J., Booth, A., 2009. A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Inf. Libr. J.* 26 (2), 91–108.
- Greenland, K., De Jager, C., Heuvelink, A., Van Der Zwaluw, K., Heck, M., Notermans, D., Van Pelt, W., Friesema, I., 2009. Nationwide outbreak of STEC O157 infection in the Netherlands, december 2008-january 2009: continuous risk of consuming raw beef products. *Euro Surveill.* 14 (8), 19129.
- Guillois, Y., Abravanel, F., Miura, N., Pavio, N., Vaillant, V., Lhomme, S., Le Guyader, F. S., Rose, N., Le Saux, J.C., King, L.A., Izopet, J., 2016. High proportion of asymptomatic infections in an outbreak of hepatitis E associated with a spit-roasted piglet, France, 2013. *Clin. Infect. Dis.* 62 (3), 351–357.
- Heaton, D., Huang, S., Shiau, R., Casillas, S., Straily, A., Kong, L.K., Ng, V., Petru, V., 2018. Trichinellosis outbreak linked to consumption of privately raised raw boar meat – California, 2017. *Morb. Mortal. Wkly Rep.* 67 (8), 247.
- Helmuth, I.G., Espenhain, L., Ethelberg, S., Jensen, T., Kjeldgaard, J., Litrup, E., Schjørring, S., Müller, L., 2019. An outbreak of monophasic *salmonella typhimurium* associated with raw pork sausage and other pork products, Denmark 2018–19. *Epidemiol. Infect.* 147 (315), 1–7.
- Hess, I.M.R., Neville, L.M., McCarthy, R., Shadbolt, C.T., McNulty, J.M., 2008. *Salmonella typhimurium* 197 outbreak linked to the consumption of lambs' liver in Sydney, NSW. *Epidemiol. Infect.* 136 (4), 461–467.
- Holzbauer, S.M., Agger, W.A., Hall, R.L., Johnson, G.M., Schmitt, D., Garvey, A., Bishop, H.S., Rivera, H., De Almeida, M.E., Hill, D., Stromberg, B.E., 2014. Outbreak of *Trichinella spiralis* infections associated with a wild boar hunted at a game farm in Iowa. *Clin. Infect. Dis.* 59 (12), 1750–1756.
- Hussain, M.A., Dawson, C.O., 2013. Economic impact of food safety outbreaks on food businesses. *Foods* 2 (4), 585–589.
- Jager, C., De Heuvelink, A., Zwaluw, K., Van Der Heck, M., Notermans, D., Pelt, W., Van Friesema, I., 2009. Nationwide outbreak of STEC O157 infection in the Netherlands, December 2008-January 2009 : continuous risk of consuming raw beef products. *Euro Surveill.* 14 (8), 7–10.
- Jay, M.T., Garrett, V., Mohle-Boetani, J.C., Barros, M., Farrar, J.A., Rios, R., Abbott, S., Sowadsky, R., Komatsu, K., Mandrell, R., Sobel, J., 2004. A multistate outbreak of *Escherichia coli* O157: H7 infection linked to consumption of beef tacos at a fast-food restaurant chain. *Clin. Infect. Dis.* 39 (1), 1–7.
- Jeffer, S.B., Kassem, I.I., Kharroubi, S.A., Abebe, G.K., 2021. Analysis of food safety management systems in the beef meat processing and distribution chain in Uganda. *Foods* 10 (10), 2244.
- Jones, G., Pihier, N., Vanbockstael, C., Le Hello, S., Cadel Six, S., Fournet, N., Jourdan-Da Silva, N., 2016. Outbreak of *Salmonella enteritidis* linked to the consumption of frozen beefburgers received from a food bank and originating from Poland: Northern France, December 2014 to April 2015. *Euro Surveill.* 21 (40).
- Jones, R.C., Reddy, V., Kornstein, L., Fernandez, J.R., Stavinsky, F., Agasan, A., Gerber, S. L., 2004. *Salmonella enterica* serotype Uganda infection in New York City and Chicago. *Emerg. Infect. Dis.* 10 (9), 1665.
- Kawakami, V., Bottichio, L., Lloyd, J., Carleton, H., Leeper, M., Olson, G., Li, Z., Kissler, B., Angelo, K.M., Whitlock, L., Sinatra, J., 2019. Multidrug-resistant *Salmonella* 14, [5], 12: i- and *Salmonella infantis* infections linked to whole roasted pigs from a single slaughter and processing facility. *J. Food Prot.* 82 (9), 1615–1624.
- Kennedy, T.G., Giotis, E.S., McKeivitt, A.I., 2014. Microbial assessment of an upward and downward dehiding technique in a commercial beef processing plant. *Meat Sci.* 97 (4), 486–489.

- King, L.A., Loukiadis, E., Mariani-Kurkdjian, P., Haeghebaert, S., Weill, F.X., Baliere, C., Ganet, S., Gouali, M., Vaillant, V., Pihier, N., Callon, H., 2014. Foodborne transmission of sorbitol-fermenting *Escherichia coli* O157: [H7] via ground beef: an outbreak in northern France, 2011. *Clin. Microbiol. Infect.* 20 (12), 1136–1144.
- King, L.A., Maillies, A., Mariani-Kurkdjian, P., Vernozzy-Rozand, C., Montet, M.P., Grimont, F., Pihier, N., Devalk, H., Perret, F., Bingen, E., Espié, E., 2009. Community-wide outbreak of *Escherichia coli* O157: H7 associated with consumption of frozen beef burgers. *Epidemiol. Infect.* 137 (6), 889–896.
- Kitchenham, B., 2004. In: *Procedures for performing systematic reviews*, 33. Keele University, Keele, UK, pp. 1–26, 2004.
- Kivi, M., Hofhuis, A., Notermans, D.W., Wannet, W.J.B., Heck, M.E.O.C., Van De Giessen, A.W., Van Duynhoven, Y.T.H.P., Stenvers, O.F.J., Bosman, A., Van Pelt, W., 2007. A beef-associated outbreak of salmonella typhimurium DT104 in the Netherlands with implications for national and international policy. *Epidemiol. Infect.* 135 (6), 890–899.
- Lachmann, R., Halbedel, S., Adler, M., Becker, N., Allerberger, F., Holzer, A., Boone, I., Falkenhörst, G., Kleta, S., Al Dahouk, S., Stark, K., 2021. Nationwide outbreak of invasive listeriosis associated with consumption of meat products in health care facilities, Germany, 2014–2019. *Clin. Microbiol. Infect.* 27 (7), 1035–e1.
- Lianou, A., Panagou, E.Z., Nychas, G.J.E., 2017. Meat safety—I foodborne pathogens other biological issues. In: *Lawrie's Meat Science*. Woodhead Publishing, pp. 521–552.
- Limon, G., Ulziibat, G., Sandag, B., Dorj, S., Purevtseren, D., Khishgee, B., Basan, G., Bandi, T., Ruuragch, S., Bruce, M., Rushton, J., 2020. Socio-economic impact of foot-and-mouth disease outbreaks and control measures: an analysis of Mongolian outbreaks in 2017. *Transbound. Emerg. Dis.* 67 (5), 2034–2049.
- Luzzi, I., Galetta, P., Massari, M., Rizzo, C., Filetici, E., Cawthorne, A., Tozzi, A., Argentieri, A., Bilei, S., Busani, L., Gnesivo, C., Pendenza, A., Piccoli, A., Napoli, P., Loffredo, R., Trinito, M.O., Santarelli, E., Atti, M.L., Dionisi, A.M., 2007. An Eastern outbreak of *Salmonella typhimurium* DT104A associated with traditional pork salami in Italy. *Euro Surveill.* 12 (4), 11–12.
- Malek, M., Barzilay, E., Kramer, A., Camp, B., Jaykus, L.A., Escudero-Abarca, B., Derrick, G., White, P., Gerba, C., Higgins, C., Vinje, J., 2009. Outbreak of norovirus infection among river rafters associated with packaged delicatessen meat, grand canyon, 2005. *Clin. Infect. Dis.* 48 (1), 31–37.
- Maurella, C., Gallina, S., Ru, G., Adriano, D., Bellio, A., Bianchi, D.M., Chiavacci, L., Crescio, M.L., Croce, M., D'Errico, V., Dupont, M.F., 2018. Outbreak of febrile gastroenteritis caused by *Listeria monocytogenes* 1/2a in sliced cold beef ham, Italy, may 2016. *Euro Surveill.* 23 (10), 17-00155.
- McEvoy, J.M., Doherty, A.M., Finnerty, M., Sheridan, J.J., McGuire, L., Blair, I.S., McDowell, D.A., Harrington, D., 2000. The relationship between hide cleanliness and bacterial numbers on beef carcasses at a commercial abattoir. *Appl. Microbiol.* 30 (5), 390–395.
- McLaughlin, J.B., Castronale, L.J., Gardner, M.J., Ahmed, R., Gessner, B.D., 2006. Outbreak of multidrug-resistant *Salmonella typhimurium* associated with ground beef served at a school potluck. *J. Food Prot.* 69 (3), 666–670.
- Mead, P.S., Dunne, E.F., Graves, L., Wiedmann, M., Patrick, M., Hunter, S., Salehi, E., Mostashari, F., Craig, A., Mshar, P., Bannerman, T., 2006. Nationwide outbreak of listeriosis due to contaminated meat. *Epidemiol. Infect.* 134 (4), 744–751.
- Messiaen, P., Forier, A., Vanderschueren, S., Theunissen, C., Nijs, J., Van Esbroeck, M., Bottaue, E., De Schrijver, K., Gyssens, I.C., Cartuyvels, R., Dorny, P., 2016. Outbreak of trichinellosis related to eating imported wild boar meat, Belgium, 2014. *Euro Surveill.* 21 (37), 30341.
- Min, M., Bai, L., Peng, X., Guo, L., Wan, K., Qiu, Z., 2021. An outbreak of botulinum types A, B, and E associated with vacuum-packaged salted fish and ham. *J. Emerg. Med.* 60 (6), 760–763.
- Møller, L.N., Petersen, E., Kapel, C.M.O., Melbye, M., Koch, A., 2005. Outbreak of trichinellosis associated with consumption of game meat in West Greenland. *Vet. Parasitol.* 132 (1–2 SPEC. ISS.), 131–136.
- Nakanwagi, M., Ario, A.R., Kwagonza, L., Aceng, F.L., Mwesigye, J., Bulage, L., Buule, J., Sendagala, J.N., Downing, R., Zhu, B.P., 2020. Outbreak of gastrointestinal anthrax following eating beef of suspicious origin: Isingiro District, Uganda, 2017. *PLoS Negl. Trop. Dis.* 14 (2), 8026.
- Needleman, I.G., 2002. A guide to systematic reviews. *J. Clin. Periodontol.* 29, 6–9.
- Nel, S., Lues, J.F.R., Buys, E.M., Venter, P., 2004. Bacterial populations associated with meat from the deboning room of a high throughput red meat abattoir. *Meat Sci.* 66 (3), 667–674.
- Nelson, M., Wright, T.L., Pierce, A., Krogwold, R.A., 2003. A common-source outbreak of trichinosis from consumption of bear meat. *J. Environ. Health* 65, 16–19.
- Noël, H., Dominguez, M., Weill, F.X., Brisabois, A., Duchazeaubeneix, C., Kerouanton, A., Delmas, G., Pihier, N., Couturier, E., 2006. Outbreak of salmonella enterica serotype Manhattan infection associated with meat products, France, 2005. *Euro Surveill.* 11 (11), 9–10.
- Nørnung, B., Andersen, J.K., Buncic, S., 2009. Main concerns of pathogenic microorganisms in meat. In: *Toldrá, F. (Ed.), Safety of Meat and Processed Meat*. Food Microbiology and Food Safety, Springer, New York, NY, pp. 3–29.
- Okpo, E., Leith, J., Smith-Palmer, A., Bell, J., Parks, D., Browning, F., Byers, L., Corrigan, H., Webster, D., Karcher, A.M., Murray, A., 2015. An outbreak of an unusual strain of *Listeria monocytogenes* infection in north-East Scotland. *J. Infect. Public Health* 8 (6), 612–618.
- Omer, M.K., Alvarez-Ordóñez, A., Prieto, M., Skjerve, E., Asehun, T., Alvsøike, O.A., 2018. A systematic review of bacterial foodborne outbreaks related to red meat and meat products. *Foodborne Pathog. Dis.* 15 (10), 598–611.
- Pichler, J., Much, P., Kasper, S., Fretz, R., Auer, B., Kathan, J., Mann, M., Huhulescu, S., Ruppitsch, W., Pietzka, A., Silberbauer, K., 2009. An outbreak of febrile gastroenteritis associated with jellied pork contaminated with *Listeria monocytogenes*. *Wien. Klin. Wochenschr.* 121 (3), 149–156.
- Plumb, I.D., Schwensohn, C.A., Gieraltowski, L., Teclé, S., Schneider, Z.D., Freiman, J., Cote, A., Noveroske, D., Kolsin, J., Brandenburg, J., Chen, J.C., 2019. Outbreak of salmonella Newport infections with decreased susceptibility to azithromycin linked to beef obtained in the United States and soft cheese obtained in Mexico - United States, 2018–2019. *Morb. Mortal. Wkly Rep.* 68 (33), 713.
- Popović-Dragonjić, L., Kocić, I., 2018. An outbreak of human trichinellosis in the village of subotina near the town of Aleksinac. *Acta Fac. Med.* 35 (2), 140–148.
- Raguenaud, M.E., Le Hello, S., Salah, S., Weill, F.X., Brisabois, A., Delmas, G., Germonneau, P., 2012. Epidemiological and microbiological investigation of a large outbreak of monophasic *Salmonella typhimurium* 4, 5, 12: i - in schools associated with imported beef in Poitiers, France, October 2010. *Euro Surveill.* 17 (40), 20289. *European Communicable Disease Bulletin*.
- Rasmussen University Learning & Library Services, 2018. *Lecture [video]*. Available from: <https://www.youtube.com/watch?v=CfUdu6ribSU&list=PLgJOrvdmou059nFj5Znx8qgToV8FknRL&index=1&t=105s>.
- Reiterova, K., Kinčėková, J., Šnábel, V., Marucci, G., Pozio, E., Dubinský, P., 2007. *Trichinella spiralis*-outbreak in the Slovak Republic. *Infection* 35 (2), 89–93.
- Rivas, L., Dupont, P.Y., Wilson, M., Rohleder, M., Gilpin, B., 2019. An outbreak of multiple genotypes of *Listeria monocytogenes* in New Zealand linked to contaminated ready-to-eat meats—a retrospective analysis using whole-genome sequencing. *Lett. Appl. Microbiol.* 69 (6), 392–398.
- Rivero-Juarez, A., Frias, M., Martinez-Peinado, A., Riscalde, M.A., Rodriguez-Cano, D., Camacho, A., García-Bocanegra, I., Cuenca-Lopez, F., Gomez-Villamandos, J.C., Rivero, A., 2017. Familial hepatitis E outbreak linked to wild boar meat consumption. *Zoonoses Public Health* 64 (7), 561–565.
- Robertson, K., Green, A., Allen, L., Ihry, T., White, P., Chen, W.S., Douris, A., Levine, J., 2016. Foodborne outbreaks reported to the US Food Safety and Inspection Service, fiscal years 2007 through 2012. *J. Food Prot.* 79 (3), 442–447.
- Robesyn, E., De Schrijver, K., Wollants, E., Top, G., Verbeeck, J., Van Ranst, M., 2009. An outbreak of hepatitis A associated with the consumption of raw beef. *J. Clin. Virol.* 44 (3), 207–210.
- Rounds, J.M., Rigdon, C.E., Muhl, L.J., Forstner, M., Danzeisen, G.T., Koziol, B.S., Taylor, C., Shaw, B.T., Short, G.L., Smith, K.E., 2012. Non-O157 Shiga toxin-producing *Escherichia coli* associated with venison. *Emerg. Infect. Dis.* 18 (2), 279.
- Ruetsch, C., Delaunay, P., Armengaud, A., Peloux-Petiot, F., Dupouy-Camet, J., Vallée, I., Polack, B., Boireau, P., Marty, P., 2016. Inadequate labeling of pork sausages prepared in Corsica causing a trichinellosis outbreak in France. *Parasite* 23 (2016).
- Sartz, L., De Jong, B., Hjertqvist, M., Plym-Forsell, L., Alsterlund, R., Löfdahl, S., Osterman, B., Ståhl, A., Eriksson, E., Hansson, H.B., Karpman, D., 2008. An outbreak of *Escherichia coli* O157: H7 infection in southern Sweden associated with consumption of fermented sausage; aspects of sausage production that increase the risk of contamination. *Epidemiol. Infect.* 136 (3), 370–380.
- Scavia, G., Ciaravino, G., Luzzi, I., Lenglet, A., Ricci, A., Barco, L., Pavan, A., Zaffanella, F., Dionisi, A.M., 2013. A multistate epidemic outbreak of salmonella goldcoast infection in humans, June 2009 to March 2010: the investigation in Italy. *Euro Surveill.* 18 (11), 20424.
- Schellenberg, R.S., Tan, B.J., Irvine, J.D., Stockdale, D.R., Gajadhar, A.A., Serhir, B., Botha, J., Armstrong, C.A., Woods, S.A., Blondeau, J.M., McNab, T.L., 2003. An outbreak of trichinellosis due to consumption of bear meat infected with *Trichinella nativa* in 2 northern Saskatchewan communities. *J. Infect. Dis.* 188 (6), 835–843.
- Schielke, A., Rabsch, W., Prager, R., Simon, S., Fruth, A., Helling, R., Schnabel, M., Sifczyk, C., Wiecek, S., Schroeder, S., Ahrens, B., 2017. Two consecutive large outbreaks of salmonella Muenchen linked to pig farming in Germany, 2013 to 2014: is something missing in our regulatory framework? *Euro Surveill.* 22 (18), 30528.
- Schimmer, B., Nygard, K., Eriksen, H.M., Lassen, J., Lindstedt, B.A., Brandal, L.T., Kapperud, G., Aavitsland, P., 2008. Outbreak of haemolytic uraemic syndrome in Norway caused by stx 2-positive *Escherichia coli* O103: H25 traced to cured mutton sausages. *BMC Infect. Dis.* 8 (1), 1–10.
- Schneider, J.L., White, P.L., Weiss, J., Norton, D., Lidgard, J., Gould, L.H., Yee, B., Vugia, D.J., Mohle-Boetani, J., 2011. Multistate outbreak of multidrug-resistant salmonella Newport infections associated with ground beef, October to December 2007. *J. Food Prot.* 74 (8), 1315–1319.
- Schroeder, S., Harries, M., Prager, R., Höfig, A., Ahrens, B., Hoffmann, L., Rabsch, W., Mertens, E., Rimek, D., 2016. A prolonged outbreak of salmonella infantis associated with pork products in Central Germany, April–October 2013. *Epidemiol. Infect.* 144 (7), 1429–1439.
- Schumacher, A.C., Elbadawi, L.I., DeSalvo, T., Straily, A., Ajzenberg, D., Letzer, D., Moldenhauer, E., Handly, T.L., Hill, D., Dardé, M.L., Pomares, C., 2021. Toxoplasmosis outbreak associated with toxoplasma gondii-contaminated venison—high attack rate, unusual clinical presentation, and atypical genotype. *Clin. Infect. Dis.* 72 (9), 1557–1565.
- Shang, X., Tonsor, G.T., 2017. Food safety recall effects across meat products and regions. *Food Policy* 69, 145–153.
- Sharma, R.K., Raghavendra, N., Mohanty, S., Tripathi, B.K., Gupta, B., Goel, A., 2014. Clinical & biochemical profile of trichinellosis outbreak in North India. *Indian J. Med. Res.* 140 (3), 414.
- Smith, A.M., Tau, N.P., Smouse, S.L., Allam, M., Ismail, A., Ramalwa, N.R., Disenyeng, B., Ngomane, M., Thomas, J., 2019. Outbreak of *Listeria monocytogenes* in South Africa, 2017–2018: laboratory activities and experiences associated with whole-genome sequencing analysis of isolates. *Foodborne Pathog. Dis.* 16 (7), 524–530.
- Smith, B., Larsson, J.T., Lisby, M., Müller, L., Madsen, S.B., Engberg, J., Bangsbo, J., Ethelberg, S., Kemp, M., 2011. Outbreak of listeriosis caused by infected beef meat

- from a meals-on-wheels delivery in Denmark 2009. *Clin. Microbiol. Infect.* 17 (1), 50–52.
- Smith-Palmer, A., Hawkins, G., Browning, L., Allison, L., Hanson, M., Bruce, R., McElhiney, J., Horne, J., 2018. Outbreak of *Escherichia coli* O157 phage type 32 linked to the consumption of venison products. *Epidemiol. Infect.* 146 (15), 1922–1927.
- Sofos, J.N., 2008. Challenges to meat safety in the 21st century. *Meat Sci.* 78 (1–2), 3–13.
- Tada, K., Suzuki, H., Sato, Y., Morishima, Y., Nagano, I., Ishioka, H., Gomi, H., 2018. Outbreak of trichinella T9 infections associated with consumption of bear meat. *Japan. Emerg. Infect. Dis.* 24 (8), 1532–1535.
- Thomas, J., Govender, N., McCarthy, K.M., Erasmus, L.K., Doyle, T.J., Allam, M., Ismail, A., Ramalwa, N., Sekwadi, P., Ntshoe, G., Shonhiwa, A., 2020. Outbreak of listeriosis in South Africa associated with processed meat. *N. Engl. J. Med.* 382 (7), 632–643.
- Torso, L.M., Voorhees, R.E., Forest, S.A., Gordon, A.Z., Silvestri, S.A., Kissler, B., Schlackman, J., Sandt, C.H., Toma, P., Bachert, J., Mertz, K.J., 2015. *Escherichia coli* O157: H7 outbreak associated with restaurant beef grinding. *J. Food Prot.* 78 (7), 1272–1279.
- Trotz-Williams, L.A., Mercer, N.J., Walters, J.M., Maki, A.M., Johnson, R.P., 2012. Pork implicated in a Shiga toxin-producing *Escherichia coli* O157:H7 outbreak in Ontario. *Canada. Can. J. Public Health* 103 (5), 322–326.
- Tsheten, T., Tshering, D., Gyem, K., Dorji, S., Wangchuk, S., Tenzin, T., Norbu, L., Jamtsho, T., 2016. An outbreak of *Aeromonas hydrophila* food poisoning in deptsang village, Samdrup Jongkhar, Bhutan, 2016. *J. Res. Health Sci.* 16 (4), 224.
- Turiac, I.A., Cappelli, M.G., Olivieri, R., Angelillis, R., Martinelli, D., Prato, R., Fortunato, F., 2017. Trichinellosis outbreak due to wild boar meat consumption in southern Italy. *Parasit. Vectors* 10 (1), 107.
- Uetani, K., Nakayama, T., Ikai, H., Yonemoto, N., Moher, D., 2009. Quality of reports on randomised controlled trials conducted in Japan: evaluation of adherence to the CONSORT statement. *Intern. Med.* 48 (5), 307–313.
- Urfer, E., Rossier, P., Mean, F., Krending, M.J., Burnens, A., Bille, J., Francioli, P., Zwahlen, A., 2000. Outbreak of salmonella braenderup gastroenteritis due to contaminated meat pies: clinical and molecular epidemiology. *Clin. Microbiol. Infect.* 6 (10), 536–542.
- Van De, N., Trung, N.V., Ha, N.H., Nga, V.T., Ha, N.M., Thuy, P.T., Chai, J.Y., 2012. An outbreak of trichinosis with molecular identification of *Trichinella* sp. in Vietnam. *Korean J. Parasitol.* 50 (4), 339.
- Vogt, R.L., Dippold, L., 2005. *Escherichia coli* O157:H7 outbreak associated with consumption of ground beef, June–July 2002. *Public Health Rep.* 120 (2), 174–178.
- Wahl, E., Rømme, S., Granum, P.E., 2013. A *Clostridium perfringens* outbreak traced to temperature-abused beef stew, Norway, 2012. *Euro Surveill.* 18 (9), 1–6.
- Whelan, J., Noel, H., Friesema, I., Hofhuis, A., de Jager, C.M., Heck, M., Heuvelink, A., van Pelt, W., 2010. National outbreak of salmonella typhimurium (Dutch) phage-type 132 in the Netherlands, October to December 2009. *Euro Surveill.* 15 (44), 19705.
- Wilson, D., Dolan, G., Aird, H., Sorrell, S., Dallman, T.J., Jenkins, C., Robertson, L., Gorton, R., 2018. Farm-to-fork investigation of an outbreak of Shiga toxin-producing *Escherichia coli* O157. *Microb. Genom.* 4 (3), 2018. <https://doi.org/10.1099/mgen.0.000160>.
- Winter, C.H., Brockmann, S.O., Sonnentag, S.R., Schaupp, T., Prager, R., Hof, H., Becker, B., Stegmann, T., Roloff, H.U., Vollrath, G., Kuhm, A.E., 2009. Prolonged hospital and community-based listeriosis outbreak caused by ready-to-eat scalded sausages. *J. Hosp. Infect.* 73 (2), 121–128.
- Wójcik, O.P., Kjelso, C., Kuhn, K.G., Müller, L., Jensen, T., Kjeldsen, M.K., Ethelberg, S., 2012. Salmonella typhimurium outbreak associated with smoked pork tenderloin in Denmark, January to March 2011. *Scand. J. Infect. Dis.* 44 (12), 903–908.
- World Health Organisation (WHO), 2022. *E. coli* fact sheet. Available from: <https://www.who.int/news-room/fact-sheets/detail/e-coli>.

# QUESTIONNAIRE

## H9(25) Part 1 of 2

### Food safety incidents in the red meat industry: A review of foodborne disease outbreaks linked to the consumption of red meat and its products, 1991 to 2021

#### INSTRUCTIONS

- Read through the article and answer the multiple-choice questions provided below.
- **Some questions may have more than one correct answer;** in which case you must mark **all** the correct answers.

#### Introduction and Study Aims

**Question 1:** Where does food contamination in the red meat industry predominantly occur?

- A:** Only on the farm before animals are transported
- B:** During processing, distribution, retail, or catering
- C:** Exclusively in the consumer's home after purchase
- D:** At border inspection points during international trade

**Question 2:** What type of hazard is associated with most of the serious food safety issues that result in consumer health problems and costly product recalls?

- A:** Chemical hazards from pesticides
- B:** Physical hazards such as metal fragments
- C:** Allergenic hazards from cross-contamination
- D:** Microbial hazards, especially bacterial pathogens

**Question 3:** What was the stated purpose of conducting this comprehensive review?

- A:** To create new, stricter laws for meat processing plants in the UK
- B:** To evaluate the critical features and consequences of global foodborne disease outbreak incidents linked to red meat
- C:** To prove that all food safety incidents result in disease outbreaks
- D:** To calculate the exact cost of food safety incidents for the United States economy

#### Methodology

**Question 4:** What methodology was employed in this study to identify relevant reports on foodborne disease outbreaks?

- A:** A series of randomized controlled trials
- B:** A "systematic search and review" methodology
- C:** A qualitative analysis of social media posts
- D:** An informal survey of food business operators

**Question 5:** What was one of the key inclusion criteria for an outbreak to be selected for this review?

- A:** The outbreak must have involved at least one death
- B:** The outbreak must have occurred in the United States or Europe
- C:** The study must have reported two or more confirmed cases of illness
- D:** The report must have been written in a language other than English

#### Transmission Settings and Food Vehicles

**Question 6:** Which transmission setting was linked to the highest number of reported foodborne outbreaks in the scientific literature reviewed?

- A:** Household settings
- B:** Foodservice establishments like restaurants
- C:** Institutional settings such as schools
- D:** Meat processing establishments

**Question 7:** During beef slaughter and processing, which operational step is identified in the literature as a primary source of carcase contamination?

- A:** The chilling and refrigeration process
- B:** The stunning of the animal before slaughter
- C:** Dehiding and skinning
- D:** The final packaging of meat cuts

**Question 8:** Which two types of red meat were most frequently associated with the outbreaks analysed in this review?

- A:** Lamb and Mutton
- B:** Bear and Wild Boar
- C:** Beef and Pork
- D:** Deer and Bison

**Question 9:** Which type of food vehicle does the article categorise "Filet américain (raw beef spread)" and "Steak tartare" as?

- A:** Ready-to-eat meat products (RTEM)
- B:** Freshly processed meat products (FPM)
- C:** Cooked meat/meal products (CMM)
- D:** Cured and fermented meat products

### **Etiologic Agents (Pathogens)**

**Question 10:** Which three pathogens were reported as the most common causes of foodborne outbreaks linked to red meat in this review?

- A:** *Listeria*, *Norovirus*, and *Trichinella*
- B:** *Campylobacter*, *Staphylococcus*, and *Yersinia*
- C:** *Hepatitis E Virus*, *Toxoplasma gondii*, and *Bacillus cereus*
- D:** *Salmonella*, *E. coli*, and *Clostridium*

**THE END**

# QUESTIONNAIRE

## H9(25) Part 2 of 2

### Food safety incidents in the red meat industry: A review of foodborne disease outbreaks linked to the consumption of red meat and its products, 1991 to 2021

#### INSTRUCTIONS

- Read through the article and answer the multiple-choice questions provided below.
- **Some questions may have more than one correct answer;** in which case you must mark **all** the correct answers.

#### Etiologic Agents (Pathogens)

**Question 1:** The findings revealed a strong association between *E. coli* outbreaks and which specific type of meat?

- A:** Pork
- B:** Mutton
- C:** Beef
- D:** Bear

**Question 2:** According to the analysis, which pathogen was responsible for the highest number of total outbreaks, illnesses, hospitalisations, and deaths?

- A:** *E. coli*
- B:** *Clostridium*
- C:** *Listeria monocytogenes*
- D:** *Salmonella*

**Question 3:** What is the key to controlling the spread of Salmonella?

- A:** Irradiating all meat products before sale
- B:** Vaccinating all food handlers
- C:** Hygienic slaughtering practices and thorough cooking and refrigeration
- D:** Using stronger chemical sanitizers on processing equipment

**Question 4:** What severe, life-threatening condition mentioned in the study is specifically associated with Shiga toxin-producing *E. coli* (STEC) infection?

- A:** Salmonellosis
- B:** Botulism
- C:** Listeriosis
- D:** Haemolytic Uraemic Syndrome (HUS)

**Question 5:** How do most *E. coli* infections get transmitted to consumers from meat products?

- A:** Through the consumption of overcooked meat
- B:** Through the consumption of raw or undercooked contaminated meat
- C:** Through contact with frozen, uncontaminated meat products
- D:** Through the inhalation of aerosolized bacteria during cooking

**Question 6:** What characteristic of *Clostridium perfringens* allows it to survive heat and other environmental stresses, making it a common cause of food poisoning?

- A:** It is a spore-forming bacteria
- B:** It produces a heat-sensitive toxin
- C:** It thrives in high-acid environments
- D:** It is exclusively found on the skin of animals

#### Contributing Factors

**Question 7:** Of all the contributing factors identified from the scientific literature, which was the most frequently reported?

- A:** Cross-contamination
- B:** Poor hygiene practices
- C:** Contamination at the source
- D:** Temperature abuse

**Question 8:** The findings indicate that foodborne outbreaks are typically caused by:

- A:** A single, isolated failure in the food chain
- B:** The actions of a single uninformed consumer
- C:** A combination of more than one contributing factor
- D:** Events that can only occur in a processing establishment

**Question 9:** Which of the following is an example of "temperature abuse"?

- A:** Storing meat in a freezer that is too cold
- B:** Serving cooked meat immediately after it reaches temperature
- C:** Improper hot holding and insufficient temperature during cooking
- D:** Using a calibrated thermometer to check food temperatures

**Question 10:** Besides temperature abuse, what other factors were identified as major contributors to foodborne outbreak incidents?

- A:** Poor marketing and labelling of meat products
- B:** High meat prices and reduced consumer spending
- C:** Poor standards of hygiene, poor GMP, and cross-contamination
- D:** The use of outdated processing machinery

#### **Conclusion and Recommendations**

**Question 11:** The conclusion emphasises that preventing outbreak incidents requires control measures applied at what point in the food system?

- A:** Only at the retail level where consumers purchase meat
- B:** Primarily on the farm before animals are sent to slaughter
- C:** Exclusively within meat processing establishments
- D:** At all stages of the food chain

**Question 12:** What important implication do the study's findings have for food safety programs and policies?

- A:** They suggest that current regulations are too strict and should be relaxed
- B:** They indicate that food safety is solely the consumer's responsibility
- C:** They prove that foodborne illness from red meat is no longer a significant problem
- D:** They can inform decisions on where regulations need to be established and indicate training needs

**Question 13:** The findings point to a shared responsibility for minimising food safety risks among which group of stakeholders?

- A:** Only government regulators and inspectors
- B:** Only the food business operators and their employees
- C:** All stakeholders, including farmers, processors, retailers, caterers, and consumers
- D:** Only international trade organizations that set standards

**Question 14:** Given that meat processing establishments are a primary origin for pathogens, what is a key recommendation to prevent future outbreaks?

- A:** Consumers should stop eating red meat entirely
- B:** All meat should be sold frozen to kill pathogens
- C:** Maintaining high hygiene standards and process controls during slaughter and processing
- D:** Closing down all small-scale butcher shops and abattoirs

**Question 15:** What specific public health action is recommended as a result of the study's findings?

- A:** Mandating that all red meat be fortified with vitamins
- B:** Reducing the legal shelf life of all fresh meat products
- C:** Enhanced and sustained public education about foodborne illness risks
- D:** Increasing the price of red meat to discourage consumption

**THE END**



**MEMBER INFO**

(Complete sections marked with an asterisk \*)

**\*FOH number**

**\*HPCSA number**

**ID number**

**Personal detail changes**

**I hereby declare that the completion of this document is my own effort without any assistance.**

SIGNED:

DATE:

This activity is accredited for **FIVE (5) CLINICAL CEU'S**. You have to achieve 70% to pass this activity

**ANSWER SHEET**

**H9(25)**

Food safety incidents in the red meat industry: A review of foodborne disease outbreaks linked to the consumption of red meat and its products, 1991 to 2021

| *Time spent on activity |   |   |   |   |   | hour |   |   |   |   | min    |    |   |   |   |   |   |  |  |  |
|-------------------------|---|---|---|---|---|------|---|---|---|---|--------|----|---|---|---|---|---|--|--|--|
| PART 1                  |   |   |   |   |   |      |   |   |   |   | PART 2 |    |   |   |   |   |   |  |  |  |
|                         | A | B | C | D | E |      | A | B | C | D | E      |    | A | B | C | D | E |  |  |  |
| 1                       |   |   |   |   |   | 1    |   |   |   |   |        | 9  |   |   |   |   |   |  |  |  |
| 2                       |   |   |   |   |   | 2    |   |   |   |   |        | 10 |   |   |   |   |   |  |  |  |
| 3                       |   |   |   |   |   | 3    |   |   |   |   |        | 11 |   |   |   |   |   |  |  |  |
| 4                       |   |   |   |   |   | 4    |   |   |   |   |        | 12 |   |   |   |   |   |  |  |  |
| 5                       |   |   |   |   |   | 5    |   |   |   |   |        | 13 |   |   |   |   |   |  |  |  |
| 6                       |   |   |   |   |   | 6    |   |   |   |   |        | 14 |   |   |   |   |   |  |  |  |
| 7                       |   |   |   |   |   | 7    |   |   |   |   |        | 15 |   |   |   |   |   |  |  |  |
| 8                       |   |   |   |   |   | 8    |   |   |   |   |        |    |   |   |   |   |   |  |  |  |
| 9                       |   |   |   |   |   |      |   |   |   |   |        |    |   |   |   |   |   |  |  |  |
| 10                      |   |   |   |   |   |      |   |   |   |   |        |    |   |   |   |   |   |  |  |  |

| How relevant was the content of this activity to your scope of practice? |                          |                          |                          |
|--|--------------------------|--------------------------|--------------------------|
|  |                          |                          |                          |
| Not at all   | Somewhat                 | Mostly                   | Extremely                |
| <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

**RETURN ANSWER SHEET TO**

**Online members**, complete your answer sheet online

CPD PORTAL: <http://foh-cpd.co.za/cpd-online/cpd-user> (Username: FOH number & Password: ID)

(Results are available immediately)

**Email or hard copy members**, complete your answer sheet online or return it to:

**WHATSAPP:** 074 230 3874 **EMAIL:** info@foh-cpd.co.za

(Results will be sent via email)

You will receive a confirmation of receipt SMS within 12-24hours, if not received please send again.

**Have a suggestion or comment?**

Send it to us on **EMAIL** or **WHATSAPP**