



FOCUS ON
HEALTH

Radiography and Clinical Technology

Self-study CPD activity

Activity Reference Number:

A9(26) 2030

Topic

The SCIEPR checklist

Article(s)

The SCIEPR checklist: A tool for standardizing chest X-ray interpretation in resource-constrained settings: A pilot study

Speciality

X-Ray


Profession(s)

Radiographer
Radiologist

You can complete any one, or all the parts of this activity

Part 1 is approved for **TWO (2) Clinical** Continuing Education Units (CEU's)

Part 2 is approved for **TWO (2) Clinical** Continuing Education Units (CEU's)



The SCIEPR checklist: A tool for standardizing chest X-ray interpretation in resource-constrained settings – A pilot study

K.M. Sethole ^{a, *}, N. Mshunqane ^b

^a Department of Radiography, Faculty of Health Sciences, University of Pretoria, South Africa

^b Department of Physiotherapy, Faculty of Health Sciences, University of Pretoria, South Africa

ARTICLE INFO

Article history:

Received 10 November 2024

Received in revised form

24 February 2025

Accepted 25 February 2025

Available online 14 March 2025

Keywords:

Checklist

Pilot study

Standardization

Pattern recognition

Perception

Content validity

ABSTRACT

Introduction: Checklists improve performance in specialized fields such as radiology. The SCIEPR (Standardization, Communication, Image Evaluation, and Pattern Recognition) checklist was developed to aid nonradiologists in interpreting chest radiographs in district hospitals with no radiologists onsite. This study aims to investigate the clinical utility of the SCIEPR checklist.

Methods: A descriptive cross-sectional pilot study included 103 participants, including 40 radiographers and 63 doctors from four district hospitals. Radiographers completed sections A and B regarding imaging protocols for chest radiographs, while doctors filled out section C for systematically searching for abnormalities. After four weeks of using the checklist, the participants completed a survey comprising 23 closed-ended and seven open-ended questions. Key measures included compliance in completing the checklist and evaluating the end-user's perceptions of the checklist.

Results: Seventy-four SCIEPR checklists were adequately completed. Sections A and B had 100 % compliance. Two items were omitted from Section C. Forty-one participants completed the survey tool (22 doctors and 19 radiographers). Participants had mixed opinions on the checklist's impact on time and workload. No item changes were suggested. Participants reported that the checklist enhanced patient care, improved service quality, reduced interpretation time, and reduced patient waiting time.

Conclusion: Following the pilot study, we refined section C of the SCIEPR checklist, improving content and face validity. The SCIEPR checklist promotes interprofessional collaboration and may reduce omission errors by standardizing imaging protocols.

Implications for practice: The SCIEPR checklist is designed to enhance collaboration between radiographers and medical doctors in chest imaging and interpretation. Its main goal is to improve the consistency and accuracy of chest X-ray interpretations, particularly in resource-limited settings with no radiologist onsite.

© 2025 The Author(s). Published by Elsevier Ltd on behalf of The College of Radiographers. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Currently, there is a universal shortage of radiologists.¹ In 2024, there were an estimated 1072 registered radiologists in SA, with only one radiologist per 100,000 people,² in a population of 63 million,³ which is little compared to Europe, where there are 13 radiologists per 100,000 people.⁴ South Africa's healthcare system faces critical challenges in diagnostics. Medical doctors regularly use chest radiographs to diagnose and monitor treatment and predict outcomes for many diseases and abnormalities, including chest

diseases. However, chest radiographs are challenging to interpret owing to overlapping clinical features and atypical presentations.⁵ Although many hospitals in South Africa perform X-rays, there are few radiologists on-site to give structured X-ray reports, and medical doctors have no structured method for interpretation.⁶

A checklist is a structured tool that categorizes items in a list format, helping comprehension and retention of information.⁷ Checklists enhance performance, prevent errors,⁸ and improve task management.^{9,10} Checklists should be developed using a systematic and comprehensive approach, particularly in specialized fields such as radiology.¹¹ Checklists may also address the existing shortage of radiologists.¹² A quantitative checklist for chest interpretation was developed by the research team and a panel of local experts, comprising radiographers, medical doctors and radiologists, using an exploratory sequential mixed method design spanning three phases. In phase 1, the researchers conducted an extensive literature review

* Corresponding author. Department of Radiography, Faculty of Health Sciences, University of Pretoria, Pretoria 0084, South Africa.

E-mail addresses: margaret.sethole@up.ac.za (K.M. Sethole), nombeko.mshunqane@up.ac.za (N. Mshunqane).

on checklists for interpreting chest X-rays.¹³ The findings of the scoping review informed the construct and content domain of the checklist. The qualitative themes were used to formulate specific, measurable indicators or items, which contributed to the development of a preliminary quantitative checklist and an accompanying survey tool. The survey tool is part of the checklist to measure the end-user's perception and response rate.

In phase 2, the researchers collaborated with a panel of local experts to finalize the checklist using the Delphi technique. The team of local experts from South Africa and key informants participated in the item-writing process and provided feedback on the relevant items on the checklist. Their input helped eliminate confusion, ambiguity, or bias that could affect the responses and helped improve the preliminary checklist's relevance, clarity, and readability. The results from the Delphi technique were presented in a joint display to integrate the theory extracted from Phase 1 and the information provided by the local experts and key informants in Phase 2. The table of specifications had all the identified constructs and items generated during operationalization to formulate the SCIEPR (Standardization, Communication, Image Evaluation, and Pattern Recognition) checklist, as seen in [Appendix A](#). The checklist aims to standardize imaging protocols, enhance interprofessional communication, and reduce omission errors when interpreting chest radiographs in hospitals without radiologists. Once the SCIEPR checklist had been developed and confirmed by a panel of experts, it was ready for pilot testing.¹⁴

This study represents the third phase of checklist development. We used a pilot-testing process to evaluate the suitability of each part of the SCIEPR checklist. Given the smaller sample sizes, the pilot study assesses clarity, aesthetics, relevance, tone, response time, and interpretation. The primary focus of this phase was to determine the clinical utility of the SCIEPR checklist.

Methods

This was a quantitative descriptive cross-sectional study. Each item of the SCIEPR checklist was accompanied by open-ended questions in a survey tool. This survey tool asked the field-testing participants to assess each item's quality and offer suggestions for improvement. The survey tool is shown in [Appendix B](#). The study was reported following the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines.¹⁵

The checklist was piloted in four district hospitals in the city of Tshwane, Gauteng Province, South Africa (SA). Hospitals A, B, C and D serve rural and urban populations, with the total number of examinations done in October 2024 as shown in [Table 1](#).

On average, each hospital performs 2500 examinations on a typical busy month. According to [Table 1](#), chest examinations represented an average of 30–35 % of all diagnostic X-ray examinations. Interpretation was limited to new cases, primarily from emergency and outpatient departments. The X-ray departments in these hospitals are open 24 h a day with no dedicated radiologists. Only new cases were assessed using the SCIEPR checklist, while follow-up chest X-ray requests were excluded due to established diagnoses. The chest radiographs are interpreted by the referring doctors.

One hundred three participants, consisting of 40 qualified radiographers and 63 registered medical doctors involved in the imaging and interpreting chest radiographs, were selected via

purposive sampling. Interested participants signed an informed consent form. A biostatistician recommended a sample size of 35 radiographers and 35 medical doctors with 80 % power, assuming a null proportion of 80 % or lower for the actual survey. The pilot study minimum sample size requirement based on the ideal effect sizes can be at least 15, 22, and 24 participants.¹⁶

Participant recruitment

Medical doctors were recruited after meeting with the heads of the departments or clinical managers at each of the four hospitals. The researchers met with the doctors to introduce the study and explain the background, problem statement, and aims. The importance of the study was discussed, and the inclusion and exclusion criteria were highlighted. The researcher demonstrated how to complete the checklist.

Radiographers were recruited after scheduling a meeting with the head of the X-ray department to introduce the study. The meetings were arranged on days when there was more staff. All explanations were done as with the medical doctors.

During the introductory meetings, all prospective participants received a packet of forms, including the three SCIEPR checklists, participant consent forms, instructions on completing the checklist, and the survey tool, all bound into a booklet. The different hospitals kept their booklets in different places. Hospitals A and B: Booklets were kept at the doctors' reporting station. Hospital C: The researchers provided each participant with a booklet. Hospital D: The booklets were kept in the X-ray department.

Data collection tools

SCIEPR checklist

The SCIEPR checklist ([Appendix A](#)), includes 35 measurable items and a back page with a chest line diagram for indicating abnormalities. The booklet included instructions on how to use the checklist. No patient demographics in terms of name, address, or date of birth were captured. Only the hospital number was captured to link the checklist to the chest radiograph.

Survey tool

We developed the survey tool with the help of a biostatistician, which was confirmed by a panel of experts who were involved in the Delphi technique during checklist development. The survey tool consisted of thirty questions, 23 closed-ended and seven open-ended ([Appendix B](#)).

Data collection

SCIEPR checklist

Data were collected over four weeks in each setting from August to December 2023. The process indicators on the checklist required a "Yes" or "No" for sections A and B and a "Normal" or "Abnormal" response for section C. In section A, the radiographer must confirm the information from the X-ray request form. "Yes" indicates that the information required to justify the examination is adequately presented, while "No" indicates a lack thereof. In section B, the radiographer indicates the position and projections performed during imaging according to the requested examination. "Yes" under image evaluation of the chest radiograph confirms adequate radiographic factor, while "No" indicates inadequacy. "N/A" means that the item did not apply to the patient, such as pregnancy status for male patients. In section C, "Normal" confirms that the

Table 1
Number of general examinations done in October 2024.

Hospital	A	B	C	D
Diagnostic examinations, including chests	2844	2242	2465	2270
Total chest examinations only	886	809	823	677

anatomical structure appeared normal regarding imaging shape, position, size, and tissue densities, and “Abnormal” was selected for structures that showed abnormal anatomy or radiographic density. Abnormal findings were not described as part of this study. Radiographers completed Sections A and B of the checklist, while medical doctors completed section C following the workflow of the radiology department. All the participants were advised to complete at least three checklists to gain experience before completing the survey. The completed SCIEPR checklists and survey tools were kept in the relevant departments and collected by the researcher.

Survey tool

All participants who used the SCIEPR checklist were invited to complete a self-administered baseline survey questionnaire to assess their perspective on the checklist’s relevance, clarity, and time implications. The participants answered questions that involved the sections they completed on the SCIEPR checklist. The survey instrument included open-ended and closed-ended questions. Section A collected demographic data, including age, gender, and professional experience. Section B featured a 23-item question survey designed to evaluate the relevance, wording, clarity, and time needed to complete the checklist for chest interpretation via a 3-point Likert scale. The 3-point Likert Scale is used to gauge the intensity of opinions, attitudes, and perceptions within a given survey context.¹⁷ It offered three choices, “positive,” “neutral or not sure,” and “negative,” making it easy for participants to express their views. Section C included seven open-ended questions intended to elicit participants’ feedback and offer suggestions for any modifications.

Data analysis

SCIEPR checklist

Data from the SCIEPR checklist were organized in an Excel spreadsheet with thirty-five items arranged in columns and participants’ responses recorded in rows. The quantitative data were subsequently analyzed to verify compliance with the predefined checklist.

Survey tool

Participants’ responses for each survey item were tabulated onto an Excel spreadsheet. The data were analyzed by an independent biostatistician, who suggested grouping items to reduce the frequency of “neutral or not sure” responses.¹⁸ Scale reliability was improved when items were grouped into themes than when they were randomly arranged.¹⁸ The items were grouped into eight constructs, namely, time, workload, length of the checklist, standardization, communication, training, image evaluation, and pattern recognition. The percentage agreements for each construct were calculated for relevance, clarity and length of time. Furthermore, the responses to the open-ended questions were analyzed using content analysis. We identified specific words, phrases, and concepts and determined the frequency of occurrence. The text was coded exactly as it appeared on the survey tool, ensuring that only words that clearly articulated the intended concepts were included. Coding was manually performed. The team was able to shed light on various factors that influenced the popularity of different checklist items, thereby enhancing the understanding of participant opinions and the overall effectiveness of the SCIEPR checklist.¹⁹

Several strategies were used to mitigate bias. Firstly, consistency in interpretations was ensured by involving multiple researchers. Two primary researchers conducted the content analysis with an

assistant researcher. Additionally, certain participants were invited to review the findings to verify that the interpretations represented their beliefs. Finally, the study’s conclusions were presented to peers for review, providing an opportunity to validate the results.

Results

SCIEPR checklist

Seventy-four SCIEPR checklists were collected, and sections A, B and C were adequately completed. Sections A and B, completed by radiographers, showed 100 % compliance. Section C, completed by medical doctors, revealed that a few participants did not evaluate abnormalities in the lung parenchyma and the diaphragm, as shown in Fig. 1.

Survey tool

Of the 103 participants recruited, only 41 (40 %) participated in the survey: 22 medical doctors and 19 radiographers (Table 2). Their ages ranged from 22 to 53 years. Work experience ranged from a few months (newly qualified interns and community service radiographers) to 22 years.

The results presented in Fig. 2 show that time and workload were not well perceived by all participants. Most participants viewed the checklist as a tool for training, image evaluations, pattern recognition, standardizing protocols, and enhancing communication.

In Section C of the survey tool, participants were asked about potential changes to the checklist items. None of the participants suggested any changes to the checklist items or format. The last question explored the perception of the participants on the use of the SCIEPR checklist; results from the open-ended questionnaire were coded and categorized into the following themes:

- 1. Reduce interpretation time: Less time for interpretation helps doctors work more efficiently.




Figure 1. Results of the SCIEPR checklist for Section B compliance.

Table 2
Demographic data.

Age in years	21–30	31–40	41–50	51 and above
Radiographers	10	8	0	1
Medical doctors	11	5	3	1
Experience in years	0 to 1	2 to 5	6 to 10	Above 10
Radiographers	6	4	5	4
Medical doctors	10	6	4	2
Gender	Male	Female		
Radiographers	8	11		
Medical doctors	9	13		

2. Increase efficiency: Streamlining processes speed up doctors' workloads.
3. Identify abnormalities faster: Enhance methods for quickly identifying abnormalities.
4. Improve image quality: Checklists aid doctors in making better evaluations and more precise diagnoses.
5. Monitor patient wait times: Checklists also track waiting times in the X-ray department, improving hospital services.

These themes suggest that the SCIEPR checklist will enhance chest diagnosis and benefit doctors and patients.

Discussion

The study found that while compliance with the SCIEPR checklist was high among radiographers, some medical doctors overlooked specific areas, such as the lung parenchyma and diaphragm. Most participants viewed the checklist positively as a training tool that enhances efficiency, improves image quality, and reduces interpretation time. No participants suggested changes to the checklist items, indicating general satisfaction with its current format.

SCIEPR checklist

The SCIEPR checklist results showed that the diaphragm and lung parenchyma areas were not assessed for abnormalities by all participants, as shown in Fig. 1. These were the only items without subheadings, which could have explained the omission. These results were presented to the panel of experts to revise and refine the SCIEPR checklist. Section C was refined by adding subheadings to clarify the diaphragm (left hemidiaphragm and right hemidiaphragm) and the lung parenchyma (left lung parenchyma and right lung parenchyma). These results suggest that users should be trained to use a systematic approach during chest interpretation.²⁰ The sequence of search steps may be modified according to individual preferences. Doctors, in particular, should be encouraged to develop a consistent, systematic, and comprehensive routine for examining the images.²¹ A familiarization period may be necessary for doctors and radiographers to get used to the added workload and time needed to complete the checklist. However, efficiency and speed will undoubtedly increase with experience²¹, and no essential items of the SCIEPR checklist should be omitted.

Interpretation is a strategy or process in which radiographers apply requisite knowledge and skills, and doctors apply perception, analysis, and synthesis skills. Radiographers need to have the requisite knowledge of anatomy and pathology, skills in radiological imaging, spatial abilities, and image manipulation skills, and be familiar with clinical information and context,²² of which are represented in sections A and B of the checklist. Perception refers to identifying radiological findings using efficient search strategies, discriminating normal from abnormal findings and pattern recognition, which is represented in section C of the SCIEPR checklist. Analysis refers to examining radiological features and synthesising the radiological and clinical findings into a conclusion²²; these two processes are not part of the checklist.

The SCIEPR checklist may improve communication and standardize imaging protocols and image evaluation criteria, representing a systematic approach to pattern recognition. Usually, the referring doctor will communicate with radiographers via an X-ray request form.²³ Section A of the SCIEPR checklist should be cross-checked against the information on the X-ray request form. Radiographers should ensure that the patient information on the request form matches that recorded on the SCIEPR checklist. If there is any missing information, the radiographer can get the information from the patient during history taking or contact the referring doctor.²⁴ Any missing information must be noted under communication in section B of the SCIEPR. Section B of the SCIEPR checklist includes another section for selecting projections and ensuring that patients are positioned according to the hospital imaging protocol for initial chest examinations. Standard projections for new patients are a posteroanterior (PA) and a lateral image in an erect position.^{25,26} Two projections should be done due to the superimposition of structures in conventional radiography. The erect position is used to demonstrate any air-fluid levels below the hemidiaphragms. *Other projections can be done, depending on the patient's condition.* Section B also incorporated image evaluation criteria carried out by a radiographer. This follows a review of the image quality regarding proper positioning, breathing status, rotation, adequate radiographic exposure, and correct anatomical side markers. Any shortcomings in these aspects will impact the diagnosis.²⁷ Medical doctors are not trained to evaluate images but radiographers are.²⁸ The listed image quality factors guide all reporting doctors on the essential factors for accurate diagnosis.

Section C of the checklist listed all anatomical structures that should be evaluated for abnormal pattern recognition on a chest




Figure 2. Item grouping response rates from the Likert scale survey (A for a positive response, B for an 'not sure', and C for a negative response).

image. The list guides the user to follow a systematic approach to search for chest abnormalities. All items listed should be evaluated to avoid omission errors.

Survey tool

Several small and medium-sized district hospitals exist in our metropolises and rural areas. These hospitals have emergency centers and outpatient departments, making them generalist facilities. In district hospitals, the clinical team consists of medical officers, clinical associates, and nurses. The extended team encompasses other allied health professionals. The approach to service delivery is collaborative and team-based, allowing for adaptation to local needs. Medical doctors are a key part of the public sector workforce, and family physicians may occasionally support the team.²⁹ District hospitals cater to most South Africans and face lower human-resourcing ratios, financial constraints, and aging infrastructure.³⁰ The lack of reporting of radiographic images can lead to ineffective patient diagnosis and mismanagement of patients.¹ There is thus an ethical obligation to find new practices to ensure patients receive quality healthcare.

Participants in our study were divided in their responses regarding the time taken to complete the checklist and the ensuing workload. A total of 15–20 minutes were needed to complete the 35-item checklist during chest interpretation. Most participants ($n = 35$) were neutral, and only a few ($n = 5$) agreed that checklists enhanced performance during high workloads and stressful conditions by freeing the mental capacity to perform essential tasks correctly and ensure a healthy approach to workload.¹⁰

Earlier studies have also reported that checklists led to longer interpretation times and lengths.³¹ In the present study, the researchers assumed this perception came from participants who used the checklist only once. The themes from the open-ended questions highlighted that using the SCIEPR checklist reduced the interpretation time, and literature has shown that interpretation time improves with experience and confidence.² The checklist is envisaged to reduce patient waiting time, enhance patient care, and improve the quality of service. The checklist also encourages a systematic approach to finding abnormal patterns in chest radiographs, reducing the evaluation time and improving service delivery. South Africa has a national policy for managing patient waiting times in outpatient departments,³² which states that the average patient waiting time for services per visit at a district hospital is estimated to be 2 h, and the total time spent by a patient per visit to a hospital, should be 3 h. Staff members at every service area must record the time of the commencement of service and exit in the respective areas as outlined in the SCIEPR checklist.³²

The participants suggested that the checklist could be helpful for training purposes. Staff in district hospitals includes medical students, newly qualified medical doctors and community service radiographers. It is imperative that radiological training is improved at all levels, but especially at the undergraduate and intern levels.² We recommend that training institutions employ the SCIEPR checklist as an Objective Structured Clinical Examination (OSCE) to prepare final-year undergraduate medical and radiography students to be efficient and apply a systematic approach to chest interpretation. OSCEs are very helpful in medical education because they allow students to practice and demonstrate clinical skills in standardized medical scenarios. The recommendation aligns with Hales and Pronovost³³ and Marcovici and Taylor³⁴, who also recommended using checklists instead of lectures or instructional videos to teach students a systematic approach to chest interpretation.

There are several reasons for the lack of proficiency in radiographic interpretation. Firstly, there are no national standards for

competency for nonspecialized doctors. Secondly, in most public facilities in South Africa, radiologists typically do not report on basic examinations. Lastly, university radiological programs vary widely, often leaving doctors to self-learn during training or afterwards.³⁵ The ongoing issue of shortage of radiologists has persisted for decades, with little hope for improvement.² Artificial intelligence learning solutions remain out of reach in low-resource settings.^{36,37} Trainers are responsible for enhancing the skills of all personnel by integrating this training tool into medical curricula and extending it to district hospitals and clinics. Notably, recurring issues related to chest radiology interpretation lack systematic solutions. Introducing the new SCIEPR checklist seeks to address this persistent challenge.

This study had several limitations. Only three hospitals, instead of four, participated in the study. The checklist was stationed differently in all settings. Optimally, the checklist should be placed in the X-ray department, considering that radiographers are the first professionals to use it. According to the pilot study results, the SCIEPR checklist was revised and refined by the panel of experts; the SCIEPR checklist is ready for a more extensive field test with a large sample size to calculate an exploratory factor analysis. Future research should explore the ability of the SCIEPR checklist to improve the accuracy of chest interpretation.

Conclusion

The pilot study enhanced the content and face validity of the SCIEPR checklist. Participants approved the SCIEPR checklist, confirming its suitability for the intended users. The results showed that the SCIEPR checklist enhanced interprofessional collaboration and had the potential to reduce omission errors by standardizing imaging protocols and systematically searching for abnormalities.

Ethics approval and consent to participate

Ethical approval for this study was obtained from Faculty of Health Science research Ethics committee in the University of Pretoria, Reference number 84/2022, and the Tshwane Research committee with reference number GP_202204_001.

Written informed consent was obtained for anonymized participant information to be published in this article.

Availability of data

Data required for this study is available from: <http://hdl.handle.net/2263/100780>.

Author contributions

KMS: Conceptualisation, Methodology, Software.
KMS and NM: Data curation, Writing- Original Draft preparation.
KMS and NM: Visualisation, Investigation.
NM: Supervision.
KMS: Software, Validation.
KMS and NM: Writing- Reviewing and Editing.

Declaration of Generative AI and AI-assisted technologies in the writing process

Not applicable.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgements

I would like to express my sincere gratitude to Dr. Cheryl Tosh for her invaluable contributions in the area of scientific editing. I also wish to acknowledge Mr. Bram Sethole for his assistance in the data collection process, as well as Ms. Kutlwano Sethole for her meticulous efforts in capturing the raw data. Additionally, I extend my

appreciation to all the radiographers and medical doctors who participated in this study, along with the Heads of Departments who coordinated the meetings in the selected hospitals. This research was part-funded by the * University Capacity Development Programme for University of Pretoria* *Grant number: A1E904.

APPENDIX A. SCIEPR checklist

Section A. check information in the request form to justify examination				
Arrival time:		Hospital or patient code:		
		yes	no	
Age:				
Biological sex:				
Gender identity:				
Patient clinical history and findings:				
Last menstrual period:				
Indication(s) for examination:				
Requested examination:				
Date of previous X-rays and other related radiological examinations				
Section B. Completed by the radiographer				
Standard procedures (tick projection and position)	Projection	Erect	Supine	Decubitus
	PA			
	Lateral			
	AP			
	Oblique			
	Other (specify)			
Communication	Write patient information discovered during imaging			
Image Evaluation of the chest	Anatomical side marker: Left /right visible on correct site	yes	no	
	Area of interest: see from above lung apices to below diaphragms			
	Alignment			
	Artifacts (preventable)			
	Artifacts (non-preventable)			
	Exposures: see air filled lungs, bony structures and heart shadow			
	Inspiration: count 9-10 posterior ribs above diaphragms on erect PA			
	Position: scapulae out of lung fields			
Rotation: sternoclavicular joints equidistant from spinous processes				
Departure time:				
Section C. completed by reporting medical doctor				
Pattern recognition	Airways:	Position of trachea	Normal	Abnormal
		Bifurcation of trachea		
		Right main bronchi		
		Left main bronchi		
	Bony structures:	clavicles		
		Scapula		
		ribs		
		Spine (check lateral)		
	Circulation and lymphatics:	Sternum (check lateral)		
		Other visible skeletal structures		
		Aortic knuckle		
		Heart size:		
	Diaphragms:	Mediastinal position		
		Mediastinal contours		
	Pleura:	Hila		
left hemi-diaphragm				
Lung parenchyma:	right hemi-diaphragm			
	Costophrenic angle			
Soft tissue:	Cardio phrenic angle			
	left lung parenchyma			
Soft tissue:	right lung parenchyma			
	Around the chest			
Soft tissue:	Breast tissue			
	neck			
Abnormal findings described (not part of this study)				

APPENDIX B. Survey tool

Section A. Demographic data collected			
	experience	gender	age
Medical doctor:			
Radiographer:			
Hospital: A, B, C, and D			
Section B. Survey questions for Phase 3 using three-point Likert scale: N=X			
1. How long did it take to complete the checklist?	≥30 min	15–20 min	≤15 min
2. Comment on the workload caused using the checklist	Increased	Right	Decreased
3. Comment on the number of items included in the checklist	many	Not sure	few
4. Comment on the items of the checklist	relevant	Not sure	irrelevant
5. Would you recommend the checklist to a colleague working in urban district hospital?	Yes	Not sure	No
6. Would you recommend the checklist to a colleague working in rural hospital?	Yes	Not sure	No
7. Would you recommend the checklist to a colleague working alone after hours?	Yes	Not sure	No
8. Would you recommend the checklist for training students and newly qualified doctors	Yes	Not sure	No
9. Would you recommend the checklist to a colleague working in hospital where there are no radiologists?	Yes	Not sure	No
10. Did the checklist improve the interprofessional communication?	Yes	Not sure	No
11. Did the checklist help you accomplish your goal of evaluating the quality of the image in terms of patient position on the chest radiograph?	Yes	Not sure	No
12. Did the checklist help you accomplish your goal of evaluating the quality of the image in terms of exposure factors on the chest radiograph?	Yes	Not sure	No
13. Did the checklist help you accomplish your goal of evaluating the quality of the image in terms of respiration phase as seen on the chest radiograph?	Yes	Not sure	No
14. Did the checklist help you accomplish your goal of evaluating the quality of the image in terms of rotation on the chest radiograph?	Yes	Not sure	No
15. Did the checklist help you accomplish your goal of evaluating the quality of the image in terms of projections done for chest interpretation PA and Lat?	Yes	Not sure	No
16. Did the checklist help you accomplish your goal of identifying abnormal patterns on the heart on the chest radiograph?	Yes	Not sure	No
17. Did the checklist help you accomplish your goal of identifying abnormal patterns on lung tissues on the chest radiograph?	Yes	Not sure	No
18. Did the checklist help you accomplish your goal of identifying abnormal patterns on diaphragms on the chest radiograph?	Yes	Not sure	No
19. Did the checklist help you accomplish your goal of identifying abnormal patterns on the pleura on the chest radiograph?	Yes	Not sure	No
20. Did the checklist help you accomplish your goal of identifying abnormal patterns on all soft tissue structures on the chest radiograph?	Yes	Not sure	No
21. Did the use of a lateral image enhance your goal of identifying abnormal patterns on the images?	Yes	Not sure	No
22. Would you recommend the use of checklist in all hospitals where radiologists are not on site?	Yes	Not sure	No
23. Did the checklist help you accomplish your goal of identifying abnormal patterns on all bony structures on the chest radiograph?	Yes	Not sure	No
Section C. Open ended questionnaire			
24. What would you remove to improve communication on the checklist?			
25. What would you add to improve image evaluation on the checklist?			
26. What would you remove to improve image evaluation on the checklist?			
27. What would you add to improve the system of searching for abnormal patterns on the checklist?			
28. What would you remove to improve the system of searching for abnormal patterns on the checklist?			
29. Overall, are you satisfied with your experience using the checklist?			
30. What is your perception in the use of the checklist in hospitals where radiologists are not on site?			

References

- Van de Venter R, ten Ham-Baloyi W. Image interpretation by radiographers in South Africa: a systematic review. *Radiography* 2019;**25**(2):178–85. <https://doi.org/10.1016/j.radi.2018.12.012>.
- Omnia health magazine. *Entering New Markets: South Africa's Healthcare Landscape: radiology South Africa faces radiologist shortage amid AI adoption*. Accessed October 2024, <https://insights.omnia-health.com/management/south-africa-faces-radiologist-shortage-amid-ai-adoption>.
- Statistics South Africa. *Mid-year population estimates*; 2024. *Statistics South Africa* 2024;**P0302**. Statistical Release P03022024. Accessed July 2024, <https://www.statssa.gov.za/publications/P0302/P03022024.pdf>.
- RCR clinical radiology workforce census 2022. Royal College of Radiologists (RCR); 2023. <https://www.rcr.ac.uk/clinical-radiology/rcrclinical-radiology-workforce-census-2022>. [Accessed 19 October 2023].
- Dreyer RG, van der Merwe CM, Nicolaou MA, Richards GA. Assessing and comparing chest radiograph interpretation in the department of internal medicine at the university of the Witwatersrand medical School, according to seniority. *Afr J Thorac Crit Care Med* 2023;**29**(1):12–7. <https://doi.org/10.7196/AJTCCM.2023.v29i1.265>.
- Sethole KM, Rudman E, Hazell LJ. Methods used by general practitioners to interpret chest radiographs at district hospitals in the city of Tshwane, South Africa. *J Med Imag Radiat Sci* 2020;**51**(2):271–9. <https://doi.org/10.1016/j.jmir.2019.12.010>.
- Sharma A, Banerjee SK, Deswal M, Pattnaik S, Sagar SS, Malini KP. A checklist to improve patient safety in general radiology. *J Adv Zool* 2023;**44**. <http://www.researchgate.net/publication/374913462>.
- Williamson Nenad M. Effects of a radiography checklist on reducing retake exposures. *J Dent Hyg* 2022;**96**(2). <https://jdh.adha.org/content/jdthyg/96/2/18.full.pdf>.
- Ely JW, Graber ML, Croskerry P. Checklists to reduce diagnostic errors. *Acad Med* 2011;**86**(3):307–13. <https://doi.org/10.1097/ACM.0b013e31820824cd>.
- Gawande A. *The Checklist Manifesto*. New York, NY: Metropolitan Books; 2009.
- Nobel JM, van Geel K, Robben SGF. Structured reporting in radiology: a systematic review to explore its potential. *Eur Radiol* 2022;**32**(4):2837–54. <https://doi.org/10.1007/s00330-021-08327-5>.
- Hales B, Terblanche M, Fowler R, Sibbald W. Development of medical checklists for improved quality of patient care. *Int J Qual Health Care* 2008;**20**(1):22–30. <https://doi.org/10.1093/intqhc/mzm062>.
- Sethole KM, Mshunqane N, Maluleke K, Kekana M, Mbonambi L. Checklists for interpreting chest radiographs: a scoping review. *J Radiol Nurs* 2023. <https://doi.org/10.1016/j.jradnu.2023.07.008>.
- Prasanna A, Bajaj G, Anil MA, Bhat JS. A Delphi survey-based development and validation of cognitive-communication assessment tool for Indian pre-schoolers. *Coda* 2023;**36**(1):e20220309. <https://doi.org/10.1590/2317-1782/20232022309>.
- Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The Strengthening the Reporting of Observational Studies in Epidemiology

- (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;**370**(9596):1453–7. [https://doi.org/10.1016/S0140-6736\(07\)61602-X](https://doi.org/10.1016/S0140-6736(07)61602-X).
16. Bujang MA, Omar ED, Foo DHP, Hon YK. Sample size determination for conducting a pilot study to assess reliability of a questionnaire. *Restor Dent Endod* 2024;**49**(1):e3. <https://doi.org/10.5395/rde.2024.49.e3>.
 17. Santoso PH, Ismail R, Suhariyono S, Setiawati FA. Comparing irt properties of different Likert scale lengths: a case from attitudinal measurement on physics education research. doi:10.21203/rs.3.rs-2648939/v1.
 18. Lam TCM, Green KE, Bordignon C. Effects of item grouping and position of the “don’t know” option on questionnaire response. *Field Methods* 2002;**14**(4): 418–32. <https://doi.org/10.1177/152582202237730>.
 19. Neuendorf KA. *The content analysis guidebook*. Sage Publications, Inc; 2017. <https://doi.org/10.4135/9781071802878>.
 20. Melingui BF, Leroy-Terquem E, Palmer M, Taguebue JV, Wachinou AP, Gaudelus J, et al. Evaluation of a short training course of chest X-ray interpretation for the diagnosis of paediatric TB. *IJTLDP Open* 2024;**1**(2):76–82. <https://doi.org/10.5588/ijtdopen.23.0484>.
 21. Geftter WB, Hatabu H. Reducing errors resulting from commonly missed chest radiography findings. *Chest* 2023;**163**(3):634–49. <https://doi.org/10.1016/j.chest.2022.12.003>.
 22. van der Gijp A, van der Schaaf MF, van der Schaaf IC, Huige JC, Ravestloot CJ, Van Schaik JP, et al. Interpretation of radiological images: towards a framework of knowledge and skills. *Adv Health Sci Educ Theory Pract* 2014;**19**(4):565–80. <https://doi.org/10.1007/s10459-013-9488-y>.
 23. Lingard L, Espin S, Whyte S, Regehr G, Baker GR, Reznick R, et al. Communication failures in the operating room: an observational classification of recurrent types and effects. *Qual Saf Health Care* 2004;**13**(5):330–4. <https://doi.org/10.1136/qhc.13.5.330>.
 24. Koirala A, Bhandari S. Are clinicians communicating adequately with radiologists through radiological requisition? A clinical audit assessing current local practice. *Nep J Radiology* 2022;**12**(1):44–8. <https://doi.org/10.3126/njr.v12i1.40089>. <https://www.nmj.com.np/nmj/index.php/nmj/article/view/195>.
 25. Pollard N, Lincoln M, Nisbet G, Penman M. Patient perceptions of communication with diagnostic radiographers. *Radiography* 2019;**25**(4):333–8. <https://doi.org/10.1016/j.radi.2019.04.002>.
 26. de Groot PM, Carter BW, Abbott GF, Wu CC. Pitfalls in chest radiographic interpretation: blind spots. In: Saunders WB, editor. *Semin roentgenol*, vol. 50; 2015. p. 197–209. <https://doi.org/10.1053/j.ro.2015.01.008>. 3.
 27. Mabotuwana T, Bhandarkar VS, Hall CS, Gunn ML. Detecting technical image quality in radiology reports. In: *AMIA Annu Symp Proc. American medical. Informatics Association*; 2018. 2018.
 28. Karera A, Engel-Hills P, Davidson F. Radiology image interpretation services in a low-resource setting: medical doctors' experiences and the potential role of radiographers. *Radiography* 2024;**30**(2):560–6. <https://doi.org/10.1016/j.radi.2024.01.009>.
 29. Mash R, Nash J. District health service delivery and the contribution of family physicians. *South Afr J Sci* 2024;**120**(11/12). <https://doi.org/10.17159/sajs.2024/18712AICLE>. Art No.18712.
 30. Mkalipi TD. *A theoretical study on primary health care services within a South African context*. 2018. <http://hdl.handle.net/10948/32193>.
 31. Sibbald M, de Bruin ABH, van Merrienboer JJG. Checklists improve experts' diagnostic decisions. *Med Educ* 2013;**47**(3):301–8. <https://doi.org/10.1111/medu.12080>.
 32. *National guideline on management of patient waiting time in clinics, community health centers, and outpatient departments of public hospitals of South Africa. National Guideline on Management of Patient Waiting Time_2024.pdf*; 2023. Accessed July 2024.
 33. Hales BM, Pronovost PJ. The checklist—a tool for error management and performance improvement. *J Crit Care* 2006;**21**(3):231–5. <https://doi.org/10.1016/j.jcrc.2006.06.002>.
 34. Marcovici PA, Taylor GA. Journal Club: structured radiology reports are more complete and more effective than unstructured reports. *AJR Am J Roentgenol* 2014;**203**(6):1265–71. <https://doi.org/10.2214/AJR.14.12636>.
 35. Irusen E. Chest radiographic interpretation—an essential competency for Africa. *Afr J Thorac Crit Care Med* 2023;**29**(1):5. <https://doi.org/10.7196/AJTCCM.2023.v29i1.892>. https://hdl.handle.net/10520/ejc-m_ajtccm_v29_n1_a2.
 36. Khalifa M, Albadawy M. AI in diagnostic imaging: revolutionising accuracy and efficiency. *Comput Methods Programs Biomed Update* 2024;**5**:100146. <https://doi.org/10.1016/j.cmpbup.2024.100146>.
 37. Hoti K, Weidmann AE. Encouraging dissemination of research on the use of artificial intelligence and related innovative technologies in clinical pharmacy practice and education: call for papers. *Int J Clin Pharm* 2024;**46**(4):777–9. <https://doi.org/10.1186/s12909-023-04698-z38>.

QUESTIONNAIRE

A9(26) Part 1 of 2

The SCIEPR checklist: A tool for standardizing chest X-ray interpretation in resource-constrained settings: A pilot study

INSTRUCTIONS

- Read through the article and answer the multiple-choice questions provided below.
- All questions have only **ONE** correct answer

Question 1: What does SCIEPR stand for?

- A:** Standardized Chest Imaging and Evaluation Protocol
- B:** Systematic Chest Interpretation and Evaluation for Radiologists
- C:** Standardization, Communication, Image Evaluation, and Pattern Recognition
- D:** Structured Chest Imaging and Pattern Recognition

Question 2: What was the main aim of the SCIEPR checklist?

- A:** To replace radiologists in urban hospitals
- B:** To aid non-radiologists in interpreting chest X-rays in district hospitals
- C:** To train radiographers in CT imaging
- D:** To automate radiology reporting

Question 3: Which professionals completed Section C of the SCIEPR checklist?

- A:** Radiographers
- B:** Nurses
- C:** Physiotherapists
- D:** Medical doctors

Question 4: What percentage of compliance was achieved in Sections A and B?

- A:** 75%
- B:** 90%
- C:** 85%
- D:** 100%

Question 5: Which anatomical areas were most commonly omitted in Section C?

- A:** Airways and pleura
- B:** Lung parenchyma and diaphragm
- C:** Soft tissue and bony structures
- D:** Circulation and heart

Question 6: What type of study was conducted to evaluate the SCIEPR checklist?

- A:** Randomized controlled trial
- B:** Longitudinal cohort study
- C:** Descriptive cross-sectional pilot study
- D:** Case-control study

Question 7: How many participants were recruited for the pilot study?

- A:** 74
- B:** 103
- C:** 41
- D:** 86

Question 8: What was the average percentage of chest X-rays among total diagnostic exams?

- A:** 10–15%
- B:** 20–25%
- C:** 30–35%
- D:** 40–45%

Question 9: What was the main reason for refining Section C of the checklist?

- A:** To add subheadings for clarity
- B:** To include patient demographics
- C:** To reduce checklist length
- D:** To improve image quality

Question 10: What tool was used to develop the checklist with expert input?

- A:** SWOT analysis
- B:** Delphi technique
- C:** Fishbone diagram
- D:** Root cause analysis

THE END

QUESTIONNAIRE

A9(26) Part 2 of 2

The SCIEPR checklist: A tool for standardizing chest X-ray interpretation in resource-constrained settings: A pilot study

INSTRUCTIONS

- Read through the article and answer the multiple-choice questions provided below.
- All questions have only **ONE** correct answer

Question 1: What was the most common perception of the checklist's impact on time and workload?

- A:** It reduced workload significantly
- B:** It eliminated errors completely
- C:** It was irrelevant
- D:** It was time-consuming

Question 2: What type of scale was used in the survey tool?

- A:** 5-point Likert scale
- B:** Binary scale
- C:** 3-point Likert scale
- D:** Visual analog scale

Question 3: Which section of the checklist includes image evaluation criteria?

- A:** Section B
- B:** Section A
- C:** Section C
- D:** Appendix B

Question 4: What is the recommended projection for new chest X-ray patients?

- A:** AP only
- B:** Lateral only
- C:** PA and lateral
- D:** Supine

Question 5: What was the average time needed to complete the checklist?

- A:** 5 minutes
- B:** 10 minutes
- C:** 30 minutes
- D:** 15-20 minutes

Question 6: What is the primary benefit of using the SCIEPR checklist in district hospitals?

- A:** Reduces radiation exposure
- B:** Improves consistency and accuracy
- C:** Increases patient volume
- D:** Replaces clinical history

Question 7: What does "Abnormal" indicate in Section C?

- A:** Normal anatomical structure
- B:** Missing patient data
- C:** Abnormal anatomy or radiographic density
- D:** Poor image quality

Question 8: Which group showed the highest compliance in completing the checklist?

- A:** Medical doctors
- B:** Radiographers
- C:** Nurses
- D:** Interns

Question 9: What is one proposed use of the SCIEPR checklist in education?

- A:** OSCE training
- B:** Lecture tool
- C:** Online quiz
- D:** Simulation software

Question 10: What national challenge does the SCIEPR checklist aim to address?

- A:** Shortage of radiologists
- B:** Poor internet access
- C:** High cost of imaging
- D:** None of the above

THE END



FOCUS ON HEALTH

www.foh-cpd.co.za

Cell: 074 230 3874

Tel: 012 653 0133 / 2373 / 2394

Mon-Fri: 08:00-16:00

400 Theuns van Niekerk Street

Wierda Park

0157

PO Box 71

Wierda Park

0149

MEMBER INFO

(Complete sections marked with an asterisk *)

*FOH number

*HPCSA number

ID number

Personal detail changes

I hereby declare that the completion of this document is my own effort without any assistance.

SIGNED:

DATE:

This activity is accredited for **FOUR (4) CLINICAL** CEU'S. You have to achieve **70%** to pass this activity.

ANSWER SHEET

A9(26)

The SCIEPR checklist: A tool for standardizing chest X-ray interpretation in resource-constrained settings: A pilot study

*Time spent on activity												hour min											
You can complete any one, or all the parts of this activity.																							
PART 1						PART 2																	
	A	B	C	D	E		A	B	C	D	E		A	B	C	D	E		A	B	C	D	E
1						6						1						6					
2						7						2						7					
3						8						3						8					
4						9						4						9					
5						10						5						10					

How relevant was the content of this activity to your scope of practice?			
			
Not at all	Somewhat	Mostly	Extremely

RETURN ANSWER SHEET TO

Online members, complete your answer sheet online

CPD PORTAL: <http://foh-cpd.co.za/cpd-online/cpd-user> (Username: FOH number & Password: ID)

(Results are available immediately)

Email or hard copy members, complete your answer sheet online or return it to:

WHATSAPP: 074 230 3874 EMAIL: info@foh-cpd.co.za

(Results will be sent via email)

You will receive a confirmation of receipt either by Whatsapp or email within 12-24hours, if not received please send again.

Have a suggestion or comment?

Send it to us on EMAIL or WHATSAPP